

# Positive operators arising from contractions

G. Geher

Let  $\mathcal{H}$  be a complex Hilbert space. If  $T \in \mathcal{B}(\mathcal{H})$  is a contraction i.e.:  $\|T\| \leq 1$ , then the sequence  $\{T^{*n}T^n\}_{n=1}^{\infty}$  of positive operators is decreasing, so it has a positive limit in the strong operator topology (SOT):

$$A_T = \lim_{n \rightarrow \infty} T^{*n}T^n.$$

We say that  $A_T$  is induced by  $T$ , or  $A_T$  is the asymptotic limit of  $T$ . The subspace

$$\mathcal{N}(A_T) = \mathcal{H}_0(T) := \{x \in \mathcal{H} : \lim_{n \rightarrow \infty} \|T^n x\| = 0\}$$

is hyperinvariant for  $T$  and it is called the stable subspace of  $T$ . The subspace

$$\mathcal{N}(A_T - I) = \mathcal{H}_1(T) := \{x \in \mathcal{H} : \lim_{n \rightarrow \infty} \|T^n x\| = \|x\|\}$$

is the largest invariant subspace where  $T$  is an isometry.

We will describe those positive operators that arises from a contraction in such a way. Moreover we can ensure uniform convergence, and expect from the case when  $0 < \dim \mathcal{H}_1 < \aleph_0$  we can choose a co-stable contraction i.e.:  $\mathcal{H}_0(T^*) = 0$ . After that we give some sufficient condition for two contraction, having the same asymptotic limit.

## References

- [C] J. B. Conway: A course in functional analysis. 2nd ed. Springer-Verlag, 1990.
- [CF] G. Cassier and T. Fack: *Contractions in Von Neumann Algebras*, Journal of Functional Analysis **135**, 297-338 (1996).
- [K1] L. Kerchy: *Unitary asymptotes and quasianalyticity*, to appear.

- [Ku] C. S. Kubrusly: An introduction to models and decompositions in operator theory, Birkhäuser, 1997.
- [NFBK] B. Sz-Nagy, C. Foias, H. Bercovici and L. Kerchy: Harmonic Analysis of Operators on Hilbert space, Springer, 2010.