

# Ergodic properties of $\beta$ -adic Halton sequences

Markus Hofer\* Maria Rita Iacò† Robert Tichy‡

## Abstract

We investigate a parametric extension of the classical  $s$ -dimensional Halton sequence where the bases are special Pisot numbers. In a one-dimensional setting the properties of such sequences have already been investigated by several authors [5, 7, 22, 27]. We use methods from ergodic theory in order to investigate the distribution behavior of multidimensional versions of such sequences. As a consequence it is shown that the Kakutani-Fibonacci transformation is uniquely ergodic.

## 1 Introduction

In this article we consider the distribution properties of deterministic point sequences in  $[0, 1)^s$ . We use the following notation: for two points  $\mathbf{a}, \mathbf{b} \in [0, 1)^s$  put  $\mathbf{a} \leq \mathbf{b}$  and  $\mathbf{a} < \mathbf{b}$  if the corresponding inequalities hold in each coordinate; furthermore, we write  $[\mathbf{a}, \mathbf{b})$  for the set  $\{\mathbf{x} \in [0, 1)^s : \mathbf{a} \leq \mathbf{x} < \mathbf{b}\}$ , and such a set is called an  $s$ -dimensional interval. Moreover we denote by  $\mathbf{1}_I$  the indicator function of the set  $I \subseteq [0, 1)^s$  and by  $\lambda_s$  the  $s$ -dimensional Lebesgue measure, we use  $\lambda$  instead of  $\lambda_1$ . Vectors are written in bold fonts and  $\mathbf{0}$  denotes the  $s$ -dimensional vector  $(0, \dots, 0)$ .

A sequence  $(\mathbf{x}_n)_{n \in \mathbb{N}}$  of points in  $[0, 1)^s$  is defined to be uniformly distributed modulo 1 (u.d.) if

$$\lim_{N \rightarrow \infty} \frac{1}{N} \sum_{n=1}^N \mathbf{1}_{[\mathbf{a}, \mathbf{b})}(\mathbf{x}_n) = \lambda_s([\mathbf{a}, \mathbf{b}))$$

for all  $s$ -dimensional intervals  $[\mathbf{a}, \mathbf{b}) \subseteq [0, 1)^s$ . A characterization of uniform distribution is due to Weyl [29]: a sequence  $(\mathbf{x}_n)_{n \in \mathbb{N}}$  of points in  $[0, 1)^s$  is u.d.

---

\*Johannes Kepler University, Institute of Financial Mathematics, Altenbergerstrasse 69, 4040 Linz, Austria. e-mail: [markus.hofer@tugraz.at](mailto:markus.hofer@tugraz.at). The author is supported by the Austrian Science Fund (FWF), Project P21943

†Graz University of Technology, Institute of Mathematics A, Steyrergasse 30, 8010 Graz, Austria and University of Calabria, Department of Mathematics and Computer Science, Via P. Bucci 30B, 87036 Arcavacata di Rende (CS), Italy. e-mail: [iaco@math.tugraz.at](mailto:iaco@math.tugraz.at) The author is partially supported by the Austrian Science Fund (FWF): W1230, Doctoral Program “Discrete Mathematics”.

‡Graz University of Technology, Institute of Mathematics A, Steyrergasse 30, 8010 Graz, Austria. e-mail: [tichy@tugraz.at](mailto:tichy@tugraz.at).

**Mathematics Subject Classification:** 11J71, 11A67, 37A05

**Keywords:** uniform distribution, low-discrepancy sequences, numeration systems

if and only if for every real-valued continuous function  $f$  the relation

$$\lim_{N \rightarrow \infty} \frac{1}{N} \sum_{n=1}^N f(\mathbf{x}_n) = \int_{[0,1]^s} f(\mathbf{x}) d\mathbf{x}$$

holds. Weyl's criterion suggests a numerical integration technique which is usually called Quasi Monte Carlo (QMC) integration. However, this characterization gives no information on the quality of the estimator.

The Koksma–Hlawka inequality [18] states that the error term of QMC integration can be bounded by the product of the variation of  $f$  (in the sense of Hardy and Krause), denoted by  $V(f)$ , and the so-called star-discrepancy  $D_N^*$  of the point sequence  $(\mathbf{x}_n)_{n \in \mathbb{N}}$ , i.e.

$$\left| \frac{1}{N} \sum_{n=1}^N f(\mathbf{x}_n) - \int_{[0,1]^s} f(\mathbf{x}) d\mathbf{x} \right| \leq V(f) D_N^*(\mathbf{x}_n),$$

where  $D_N^*$  is defined by

$$D_N^*(\mathbf{x}_n) = D_N^*(\mathbf{x}_1, \dots, \mathbf{x}_N) = \sup_{\mathbf{a} \in [0,1]^s} \left| \frac{1}{N} \sum_{n=1}^N \mathbf{1}_{[\mathbf{0}, \mathbf{a}]}(\mathbf{x}_n) - \lambda_s([\mathbf{0}, \mathbf{a}]) \right|.$$

Thus in order to minimize the integration error we have to use point sequences with small discrepancy. There are several constructions for sequences which achieve a star-discrepancy of order  $\mathcal{O}(N^{-1}(\log N)^s)$ , so-called low-discrepancy sequences. This convergence rate, which is best possible among known sequences, is for all  $s \geq 1$  better than that of the probabilistic error of the standard Monte Carlo method, where a sequence of random instead of deterministic points is used. QMC integration is successfully used in several different areas of applied mathematics, for example in actuarial and financial mathematics, where high-dimensional numerical integrals frequently appear, see e.g. [4, 23]. For a detailed survey on low-discrepancy sequences and their applications we refer to [10].

In this article we construct point sequences by a combination of methods from uniform distribution theory and dynamical systems. In particular we give answers to problems considered in [1] and [8]. For the basic definitions we refer to these articles and to standard text books such as [28].

An important tool for analyzing the properties of a dynamical system is the construction of an isometry to a known ergodic or uniquely ergodic system. A classical example of a uniquely ergodic system is  $(\mathbb{Z}_b, \tau)$  (see for instance, [28]), where  $\mathbb{Z}_b$  is the compact group of  $b$ -adic integers and  $\tau : \mathbb{Z}_b \rightarrow \mathbb{Z}_b$  the addition-by-one map (called odometer). In the sequel we shortly recall the connection between  $(\mathbb{Z}_b, \tau)$  and low-discrepancy sequences on  $[0, 1)$ . For an integer  $b \geq 2$ , every  $z \in \mathbb{Z}_b$  has a unique expansion of the form

$$z = \sum_{j \geq 0} z_j b^j$$

with digits  $z_j \in \{0, 1, \dots, b-1\}$ . For  $z \in \mathbb{Z}_b$  we define the  $b$ -adic Monna map  $\varphi_b : \mathbb{Z}_b \rightarrow [0, 1)$ , see also [13], by

$$\varphi_b \left( \sum_{j \geq 0} z_j b^j \right) = \sum_{j \geq 0} z_j b^{-j-1}.$$

The restriction of  $\varphi_b$  to  $\mathbb{N}_0$  is called radical-inverse function in base  $b$  and the sequence

$$(\varphi_b(n))_{n \in \mathbb{N}}$$

is the so-called van der Corput sequence in base  $b$  which is a low-discrepancy sequence.

The Monna map is continuous and surjective but not injective. In order to make it an isomorphism we only consider the so-called regular representations, i.e. representations with infinitely many digits  $z_j$  different from  $b-1$ . The Monna map restricted to regular representations admits an inverse (called pseudo-inverse)  $\varphi_b^{-1} : [0, 1) \rightarrow \mathbb{Z}_b$ , defined by

$$\varphi_b^{-1} \left( \sum_{j \geq 0} z_j b^{-j-1} \right) = \sum_{j \geq 0} z_j b^j,$$

where  $\sum_{j \geq 0} z_j b^{-j-1}$  is a  $b$ -adic rational in  $[0, 1)$ .

Moreover  $\varphi_b$  is measure preserving from  $\mathbb{Z}_b$  onto  $[0, 1)$  and it transports the normalized Haar measure on  $\mathbb{Z}_b$  to the Lebesgue measure. Hence, by the unique ergodicity of  $\tau$  it follows that the sequence  $(\tau^n z)_{n \in \mathbb{N}}$  is uniformly distributed in  $\mathbb{Z}_b$  for all  $z \in \mathbb{Z}_b$ , in particular for  $z = 0$ . Thus the van der Corput sequence  $(\varphi_b(\tau^n 0))_{n \in \mathbb{N}}$  in base  $b$  is uniformly distributed modulo 1.

In order to construct multidimensional sequences we need a criterion to ensure that the Cartesian product of ergodic systems is again ergodic, see e.g. [13].

**Theorem 1.** *Let  $\mathcal{T}_i = (X_i, T_i)$ ,  $i = 1, \dots, s$ , be ergodic dynamical systems. Then the dynamical system  $\mathcal{T}_1 \times \dots \times \mathcal{T}_s$  is ergodic if and only if for all  $i, j \in \{1, \dots, s\}$ ,  $i \neq j$ , the discrete parts of the spectra of  $T_i$  and  $T_j$  intersect only at 1.*

In [13], the spectrum of  $(\mathbb{Z}_b, \tau)$  is given explicitly. Furthermore the authors show that if  $b_1, b_2$  denote positive integers and  $\tau_{b_i}$  denotes the addition-by-one on  $\mathbb{Z}_{b_i}$ , then the dynamical systems  $(\mathbb{Z}_{b_1}, \tau_{b_1})$  and  $(\mathbb{Z}_{b_2}, \tau_{b_2})$  are spectrally disjoint if and only if  $b_1$  and  $b_2$  are coprime. This is exactly the condition proved by Halton in [17] in order to obtain a low-discrepancy sequence in  $[0, 1)^s$  by combining coordinatewise van der Corput sequences. The resulting sequence  $(\phi_{\mathbf{b}}(n))_{n \in \mathbb{N}} = (\phi_{b_1}(n), \dots, \phi_{b_d}(n))_{n \in \mathbb{N}}$  is called  $\mathbf{b}$ -adic Halton sequence, where  $\mathbf{b}$  is an  $s$ -dimensional vector of pairwise coprime integers  $b_i$ ,  $i = 1, \dots, s$ .

The aim of the present article is to extend the above idea to point sequences with irrational bases. In a one-dimensional setting such sequences were investigated by several authors. For instance, Barat and Grabner [5] consider the so-called

$\beta$ -adic van der Corput sequence  $(\phi_\beta(n))_{n \in \mathbb{N}}$  on  $[0, 1)$  and similar constructions. They prove that  $(\phi_\beta(n))_{n \in \mathbb{N}}$  is low-discrepancy, if  $\beta$  is the characteristic root of a special linear recurrence. Ninomiya [22] considers the discrepancy of point sequences on  $[0, 1)$  for a slightly greater class of irrational bases  $\beta$ . Furthermore Steiner [26] investigates so-called bounded remainder sets in a similar setting. In a second article Steiner [27] considers van der Corput sequences on abstract numeration systems and gives conditions under which they are low-discrepancy. Discrepancy bounds for a multidimensional extension of [22] are formulated by Mori and Mori [21] by using algebraic methods. Note that the construction in [21] is different from that in the present article.

Carbone [7] and Drmota and Infusino [9] investigate the discrepancy of point sequences generated by the so-called Kakutani splitting procedure. Carbone completely characterizes the growth order of the discrepancy for a two parametric subfamily, so-called LS-sequences. Moreover, Aistleitner, Hofer and Ziegler [1] give conditions under which an  $s$ -dimensional vector of LS-sequences is not u.d. in  $[0, 1)^s$ .

The remainder of this article is structured as follows: in the next section we formulate a characterization of uniquely ergodic systems which are constructed as Cartesian products of odometers on numeration systems related to linear recurrences. In the third section we give conditions under which the  $s$ -dimensional  $\beta$ -adic Halton sequence is uniformly distributed sequence in  $[0, 1)^s$ . Furthermore we present a parametric class of sequences which satisfies these conditions. Finally we prove that the ergodic Kakutani-Fibonacci-transformation, presented in [8], is in fact uniquely ergodic.

## 2 General $G$ -odometers

In this section we consider odometers on numeration systems with respect to a linear recurring base sequence. For a detailed discussion of such number systems we refer to [11, 14, 15, 16].

**Definition 1.** *Let  $(G_n)_{n \geq 0}$  be an increasing sequence of positive integers with  $G_0 = 1$ . Then every positive integer can be expanded in the following way*

$$n = \sum_{k=0}^{\infty} \varepsilon_k(n) G_k, \quad \forall n \in \mathbb{N}$$

where  $\varepsilon_k(n) \in \{0, \dots, \lfloor G_{k+1}/G_k \rfloor\}$  and  $\lfloor x \rfloor$  denotes the integral part of  $x$ , that is the greatest integer less than or equal to  $x \in \mathbb{R}$ . This expansion (called  $G$ -expansion) is uniquely determined and finite, provided that for every  $K$

$$\sum_{k=0}^{K-1} \varepsilon_k(n) G_k < G_K. \quad (1)$$

For short we write  $\varepsilon_k$  for the  $k$ -th digit of the  $G$ -expansion.

The digits  $\varepsilon_k$  can be computed by the greedy algorithm (see for instance [11]) and  $G = (G_n)_{n \geq 0}$  is called numeration system.

We denote by  $\mathcal{K}_G$  the subset of sequences that satisfy (1), and the elements of  $\mathcal{K}_G$  are called  $G$ -admissible. In order to extend the addition-by-one map defined on  $\mathbb{N}$  to  $\mathcal{K}_G$  we introduce  $\mathcal{K}_G^0 \subseteq \mathcal{K}_G$

$$\mathcal{K}_G^0 = \left\{ x \in \mathcal{K}_G : \exists M_x, \forall j \geq M_x, \sum_{k=0}^j \varepsilon_k G_k < G_{j+1} - 1 \right\}. \quad (2)$$

Put  $x(j) = \sum_{k=0}^j \varepsilon_k G_k$ , and set

$$\tau(x) = (\varepsilon_0(x(j) + 1) \dots \varepsilon_j(x(j) + 1)) \varepsilon_{j+1}(x) \varepsilon_{j+2}(x) \dots, \quad (3)$$

for every  $x \in \mathcal{K}_G^0$  and  $j \geq M_x$ . This definition does not depend on the choice of  $j \geq M_x$  and can be easily extended to sequences  $x$  in  $\mathcal{K}_G \setminus \mathcal{K}_G^0$  by  $\tau(x) = 0 = (0^\infty)$ . In this way the transformation  $\tau$  is defined on  $\mathcal{K}_G$  and it is called  $G$ -odometer. We refer to [14] for a complete survey on odometers related to general numeration systems.

In this article we consider only numeration systems where the base sequence is a linear recurrence. Let  $G_0 = 1$  and  $G_k = a_0 G_{k-1} + \dots + a_{k-1} G_0 + 1$  for  $k < d$ . Then  $G_n$  for  $n \geq d$  is determined by a recurrence of order  $d \geq 1$ , i.e.

$$G_{n+d} = a_0 G_{n+d-1} + \dots + a_{d-1} G_n, \quad n \geq 0. \quad (4)$$

The solution of the characteristic equation of the numeration system  $G$

$$x^d = a_0 x^{d-1} + \dots + a_{d-1} \quad (5)$$

plays a crucial role. We are only interested in numeration systems where the solution of (5) is a Pisot number  $\beta$ . Note that  $\beta$  is always a Pisot number if

$$a_0 \geq \dots \geq a_{d-1} \geq 1, \quad (6)$$

see [12, Theorem 2]. Parry [24] shows that in this case the so-called Parry's  $\beta$ -expansion of  $\beta$  is finite, i.e.

$$\beta = a_0 + \frac{a_1}{\beta} + \dots + \frac{a_{d-1}}{\beta^{d-1}}, \quad (7)$$

where  $a_0 = \lfloor \beta \rfloor$ . At the end of the last section we also consider numeration systems where (6) does not hold.

For numeration systems where the characteristic root  $\beta$  is a Pisot number which satisfies (7), a sum  $\sum_{k=0}^M \varepsilon_k G_k$ ,  $M < \infty$  is the expansion of an integer if and only if the digits  $\varepsilon_k$  of the  $G$ -expansion satisfy

$$(\varepsilon_k, \varepsilon_{k-1}, \dots, \varepsilon_0, 0^\infty) < (a_0, a_1, \dots, a_{d-1})^\infty, \quad (8)$$

for every  $k$  and  $<$  denoting the lexicographic order (see [24]). Representations  $(\varepsilon_k, \dots, \varepsilon_0)$  satisfying this condition are called admissible representations and thus they belong to  $\mathcal{K}_G$ .

In [14, Theorem 5], the authors show that the odometer on an admissible numeration system  $G$  is uniquely ergodic and the corresponding invariant measure  $\mu$  is given by

$$\mu(Z) = \frac{F_{K,0}\beta^{d-1} + (F_{K,1} - a_0F_{K,0})\beta^{d-2} + \dots + (F_{K,d-1} - a_0F_{K,d-2} - \dots - a_{d-2}F_{K,0})}{\beta^K(\beta^{d-1} + \beta^{d-2} + \dots + 1)}, \quad (9)$$

where  $F_{k,r} := \#\{n < G_{k+r} : (\varepsilon_0(n), \varepsilon_1(n), \dots) \in Z\}$  and  $Z$  is a cylinder with length  $k$ . Note that the corresponding formula in [14, Theorem 5] contains a misprint which was corrected in [5].

In the sequel we want to apply Theorem 1, thus we need information on the spectrum of the  $G$ -odometer. We introduce the following two hypotheses:

**Hypothesis 1** (Grabner, Tichy and Liardet [14]). *There exists an integer  $b > 0$  such that for all  $k$  and*

$$N = \sum_{i=0}^k \epsilon_i(N)G_i + \sum_{j=k+b+2}^{\infty} \epsilon_j(N)G_j,$$

*the addition of  $G_m$  to  $N$ , where  $m \geq k + b + 2$ , does not change the first  $k + 1$  digits in the greedy representation i.e.*

$$N + G_m = \sum_{i=0}^k \epsilon_i(N)G_i + \sum_{j=k+1}^{\infty} \epsilon_j(N + G_m)G_j.$$

**Hypothesis 2** (Frougny and Solomyak [12]). *The characteristic root  $\beta$  of the numeration system  $G$  is a Pisot number such that all elements of the set  $\mathbb{Z}[\beta^{-1}]$  have finite  $\beta$ -expansions.*

There are only a few results concerning Hypothesis 1. In [14] the authors remark that the Multinacci sequence, i.e.  $a_0 = \dots = a_{d-1} = 1$ , fulfills Hypothesis 1. Furthermore Bruin et al. [6] show that the numeration system with coefficients  $(a_0, a_1, a_2) = (1, 0, 1)$  fulfills Hypothesis 1.

Several researchers worked on algebraic characterizations of Pisot numbers  $\beta$  which satisfy Hypothesis 2. Frougny and Solomyak [12] show that (6) implies Hypothesis 2 and they give a full characterization of all Pisot numbers of degree two with this property. Furthermore, Hollander [19] states another sufficient condition for Hypothesis 2 and Akiyama [2] characterizes all Pisot units of degree three satisfying Hypothesis 2. Further progress was also made by Akiyama et al. [3] who prove Hypothesis 2 for a large class of Pisot numbers of degree three by using the theory of shift radix systems. Nevertheless there exists no complete algebraic characterization for Pisot numbers satisfying Hypothesis 2 of degree greater than two. Note that both hypotheses can be satisfied by the same numeration system but, to the best of the authors knowledge, it is unknown if the two hypotheses are equivalent, see [14].

Grabner, Liardet and Tichy [14, Theorem 6] and Solomyak [25, Theorem 4.1] show that the odometer on the base system  $G$  has purely discrete spectrum

provided that one of the above Hypotheses holds. Furthermore we obtain in both cases that the set of eigenvalues of the transformation is given by

$$\Gamma := \{z \in \mathbb{C} : \lim_{n \rightarrow \infty} z^{G_n} = 1\}. \quad (10)$$

**Theorem 2.** *Let  $G^1, \dots, G^s$  be numeration systems given by (4) and let the coefficients of the linear recurrences be given as  $a_j^i = b_i$ ,  $j = 0, \dots, (d_i - 1)$ ,  $i = 1, \dots, s$ , with pairwise coprime positive integers  $b_i$ ,  $i = 1, \dots, s$ . Furthermore let  $\frac{\beta_i^k}{\beta_j^l} \notin \mathbb{Q}$ , for all  $l, k \in \mathbb{N}$ , where  $\beta_1, \dots, \beta_s$  denote the characteristic roots of the numerations systems. Then the dynamical system which is constructed as the  $s$ -dimensional Cartesian product of the odometers, i.e.  $((\mathcal{K}_{G^1}, \tau_1) \times \dots \times (\mathcal{K}_{G^s}, \tau_s))$ , is uniquely ergodic.*

*Proof.* It follows by [25, Main Theorem] that each numeration system  $G^j$ ,  $j = 1, \dots, s$  fulfills Hypothesis 2, thus the components of the  $s$ -dimensional dynamical system are uniquely ergodic. Furthermore we obtain that their spectrum is given by (10). By Theorem 1, we derive that the Cartesian product is uniquely ergodic if and only if  $\Gamma_j \cap \Gamma_k = \{1\}$  for all  $1 \leq j < k \leq s$ . As noted in [14], we have

$$\lim_{n \rightarrow \infty} \frac{G_n^j}{\beta_j^n} = C_j, \quad (11)$$

where the constant  $C_j$  can be computed by residue calculus. Using the standard notation  $\sim$  for asymptotic equality (if  $n \rightarrow \infty$ ) we obtain for fixed  $l \in \mathbb{N}$

$$\begin{aligned} \exp\left(2\pi i \frac{G_n^j}{\beta_j^l}\right) &\sim \exp(2\pi i C_j \beta_j^{n-l}) \\ &\sim \exp(2\pi i G_{n-l}^j), \end{aligned}$$

and thus

$$\lim_{n \rightarrow \infty} \exp\left(2\pi i \frac{G_n^j}{\beta_j^l}\right) = \lim_{n \rightarrow \infty} \exp(2\pi i G_{n-l}^j) = 1,$$

where  $C_j$  is given by (11). Furthermore, it is easy to see that for every  $k \in \mathbb{N}$  there exists a  $n_0$  with  $b_j^k \mid G_n^j$  for all  $n \geq n_0$  and there exist no  $b', n'_0 \in \mathbb{N}$  with  $\gcd(b', b_j) = 1$  such that  $b' \mid G_n^j$  for all  $n \geq n'_0$ . Then  $\Gamma_j$  can be written as

$$\Gamma_j = \left\{ \exp\left(2\pi i \frac{c}{b_j^m \beta_j^l}\right) : m, l, c \in \mathbb{N} \cup \{0\} \right\}.$$

By Theorem 1 the  $s$ -dimensional dynamical system is ergodic. Since, by our assumption, the spectrum is purely discrete the system is isomorphic to a group rotation, and thus uniquely ergodic.  $\square$

### 3 Uniform distribution of the $\beta$ -adic Halton sequence

First we extend the definition of the Monna map to irrational bases  $\beta > 1$ . Let

$$n = \sum_{j \geq 0} \epsilon_j(n) G_j$$

be the  $G$ -expansion of an integer  $n$ . For short we write  $\epsilon_j$  and define the  $\beta$ -adic Monna map  $\phi_\beta: \mathcal{K}_G \rightarrow \mathbb{R}^+$  as

$$\phi_\beta(n) = \phi_\beta \left( \sum_{j \geq 0} \epsilon_j(n) G_j \right) = \sum_{j \geq 0} \epsilon_j(n) \beta^{-j-1}.$$

Furthermore, we define the radical inverse function as restriction of  $\phi_\beta$  on  $\mathcal{K}_G^0$  and the pseudo-inverse  $\phi_\beta^{-1}$  similarly. In this context the  $\beta$ -adic Halton sequence is given as  $(\phi_\beta(n))_{n \in \mathbb{N}} = (\phi_{\beta_1}(n), \dots, \phi_{\beta_s}(n))_{n \in \mathbb{N}}$ , where  $\beta = (\beta_1, \dots, \beta_s)$  and the  $\beta_i$  are characteristic roots of the numeration systems  $G^i$ , respectively.

Note that even if one of the Hypotheses 1 or 2 holds, this does not imply that the image of  $\mathcal{K}_G^0$  under  $\phi_\beta$  is contained and dense in  $[0, 1)$ .

**Lemma 1.** *Let  $\mathbf{a} = (a_0, \dots, a_{d-1})$ , where the integers  $a_0, \dots, a_{d-1} \geq 0$  are the coefficients defining the numeration system  $G$  and assume that the corresponding characteristic root  $\beta$  satisfies (7). Furthermore, assume that there is no  $\mathbf{b} = (b_0, \dots, b_{k-1})$  with  $k < d$  such that  $\beta$  is the characteristic root of the polynomial defined by  $\mathbf{b}$ . Then  $\phi_\beta(\mathbb{N}) \subset [0, 1)$  and  $\phi_\beta(\mathbb{N}) \not\subset [0, x)$  for all  $0 < x < 1$  if and only if  $\mathbf{a}$  can be written either as*

$$\mathbf{a} = (a_0, \dots, a_0) \text{ or as} \tag{12}$$

$$\mathbf{a} = (a_0, a_0 - 1, \dots, a_0 - 1, a_0), \tag{13}$$

where  $a_0 > 0$ .

*Proof.* It follows by (7) that

$$\frac{a_0}{\beta} + \dots + \frac{a_{d-1}}{\beta^d} = 1. \tag{14}$$

Furthermore for all admissible representations  $(\varepsilon_0, \varepsilon_1, \varepsilon_2, \dots)$  we have

$$(\varepsilon_k, \varepsilon_{k-1}, \dots, \varepsilon_0, 0^\infty) < (a_0, a_1, \dots, a_{d-1})^\infty \tag{15}$$

for every  $k$  and  $<$  denoting the lexicographic order.

We consider a representation given by  $\mathbf{c} = (c_0, \dots, c_{k-1})^\infty$  for  $k > 0$  and assume that there are no positive integers  $c'_i$  and  $m < k$  such that  $\mathbf{c} = (c'_0, \dots, c'_{m-1})^\infty$ . We obtain

$$\begin{aligned} \phi_\beta(\mathbf{c}) &= \sum_{i=0}^{\infty} \left( \frac{c_0}{\beta} + \dots + \frac{c_{k-1}}{\beta^k} \right) \left( \frac{1}{\beta^k} \right)^i \\ &= \left( \frac{c_0}{\beta} + \dots + \frac{c_{k-1}}{\beta^k} \right) \frac{1}{1 - \frac{1}{\beta^k}} \\ &= \left( \frac{c_0}{\beta} + \dots + \frac{c_{k-1}}{\beta^k} \right) \frac{\beta^k}{\beta^k - 1}. \end{aligned}$$

To finish the proof of the lemma, the following will be shown: the maximum of  $\phi_\beta(\mathbf{c})$  (extended over all representations  $\mathbf{c}$ ) is one, provided that (12) or (13) are satisfied, i.e.

$$\frac{c_0}{\beta} + \dots + \frac{c_{k-2}}{\beta^{k-1}} + \frac{c_{k-1} + 1}{\beta^k} = 1. \tag{16}$$



We obtain  $k \geq d$  since  $\mathbf{a}$  is by assumption the minimal representation of  $\beta$ . In the case  $k = d$  we get

$$\frac{c_0}{\beta} + \dots + \frac{c_{d-2}}{\beta^{d-1}} + \frac{c_{d-1} + 1}{\beta^d} = \frac{a_0}{\beta} + \dots + \frac{a_{d-1}}{\beta^d} = 1,$$

thus  $c_0 = a_0, \dots, c_{d-2} = a_{d-2}$  and  $c_{d-1} + 1 = a_{d-1}$ . Moreover, by (15), we have  $\max_{i=0, \dots, d-1} c_i \leq a_0$  and thus  $a_0 = \max(\max_{i=0, \dots, d-2} a_i, a_{d-1} - 1)$ . Let  $m \geq 0$  be the maximal integer such that  $a_i = a_0$  for  $i \leq m$ . Then, if  $a_{d-1} = a_0 + 1$  we obtain that  $\mathbf{c} = (a_0, a_1, \dots, a_{d-2}, a_0)^\infty$  contains  $m + 1$  successive  $a_0$ 's, thus  $\mathbf{c}$  is not admissible and  $a_0 = \max_{i=0, \dots, d-1} a_i$ . Similar arguments yield  $k \leq d$  since by assumption  $\mathbf{c} \neq (c'_0, \dots, c'_{m-1})^\infty$  for  $m < k$ .

Assume now that  $k = d$ ,  $a_0 = \max_{i=0, \dots, d-1} a_i$  and let  $m \geq 0$  be defined as above. By (14) and (15)  $\mathbf{c}^* = (a_0, \dots, a_m, (a_0 - 1, a_0, \dots, a_m)^\infty)$  is admissible and

$$\phi_\beta(\mathbf{c}^*) \geq 1,$$

where equality holds if  $m$  is either 0 or  $d - 1$  which are the cases (12) and (13). This yields assertion (16), thus the proof of the lemma is complete.  $\square$

If we drop the condition that  $d$  has to be minimal, we obtain the following additional cases for which the above lemma is satisfied:

$$\mathbf{a} = (a_0, \dots, a_0, a_0 + 1) \tag{17}$$

and

$$\mathbf{a} = (\mathbf{a}', \dots, \mathbf{a}', \mathbf{a}''), \tag{18}$$

where  $a_0 > 0$ ,  $\mathbf{a}', \mathbf{a}''$  are of equal length and of the form

$$\begin{aligned} \mathbf{a}' &= (a_0, \dots, a_0, a_0 - 1), & \mathbf{a}'' &= (a_0, \dots, a_0) \text{ or} \\ \mathbf{a}' &= (a_0, a_0 - 1, \dots, a_0 - 1), & \mathbf{a}'' &= (a_0, a_0 - 1, \dots, a_0 - 1, a_0). \end{aligned}$$

Note that (17) is another way to represent the  $(a_0 + 1)$ -adic number system, which is a special case of (12) and obviously fulfills the lemma. Furthermore, condition (18) is a reformulation of (12) and (13), thus in the sequel we only consider numeration systems which satisfy (12) or (13).

**Lemma 2.** *Let  $G$  be a numeration system of the form (4), assume that the coefficients of the linear recurrence are given by  $a_j = a$ ,  $j = 0, \dots, (d - 1)$ , for a positive integer  $a$  and let  $\beta$  denote the corresponding characteristic root. Then  $\mu(Z) = \lambda(\phi_\beta(Z))$  for every cylinder set  $Z$ .*

*Proof.* Let the cylinder set  $Z$  be defined by the fixed digits  $\epsilon_0, \dots, \epsilon_{k-1}$ . Assume first that  $\epsilon_{k-1} < a$ , then  $F_{k+r} = (a + 1)^r$  for  $0 \leq r < d$ . Thus, by (9), we obtain that

$$\mu(Z) = \beta^{-k}.$$

Consider the  $\beta$ -adic Monna map of  $n \in \mathbb{N}$ , i.e.

$$\phi_\beta(n) = \sum_{i=0}^{\infty} \frac{\epsilon_i}{\beta^{i+1}}.$$

If  $\epsilon_{k-1} < a$  we easily see that  $\phi_\beta(Z)$  is dense in

$$I = \left[ \sum_{i=0}^{k-1} \frac{\epsilon_i}{\beta^{i+1}}, \sum_{i=0}^{k-2} \frac{\epsilon_i}{\beta^{i+1}} + \frac{(\epsilon_{k-1} + 1)}{\beta^k} \right)$$

and that  $\phi_\beta(x') \notin I$  if  $x' \notin Z$ . Thus  $\phi_\beta(Z)$  is  $\lambda$ -measurable and  $\lambda(\phi_\beta(Z)) = \lambda(I) = \beta^{-k}$ .

Assume now that  $Z$  is defined by the fixed digits  $\epsilon_0, \dots, \epsilon_{k-2}$  and  $\epsilon_{k-1} = a$ . By the above argument we derive that a cylinder with fixed digits  $\epsilon_0, \dots, \epsilon_{k-2}$ , with  $\epsilon_{k-2} < a$ , has measure  $\beta^{-(k-1)}$ .

Now compute the measure of  $Z$ :

$$\mu(Z) = \beta^{-(k-1)} - (a-1)\beta^{-k}.$$

Next we consider  $\phi_\beta(Z)$ , hence

$$\phi_\beta(Z) = \left[ \sum_{i=0}^{k-1} \frac{\epsilon_i}{\beta^{i+1}}, \sum_{i=0}^{k-3} \frac{\epsilon_i}{\beta^{i+1}} + \frac{(\epsilon_{k-2} + 1)}{\beta^k} \right)$$

and thus  $\lambda(\phi_\beta(Z)) = \mu(Z)$ .

Let  $2 \leq h \leq \min(k, d-1)$  and consider a cylinder set  $Z$  with fixed digits  $\epsilon_0, \dots, \epsilon_{k-h-1} < a$  and  $\epsilon_{k-l} = a$  for  $l = 1, \dots, h$ . Then, the cylinder with fixed digits  $\epsilon_0, \dots, \epsilon_{k-h-1}$  has measure  $\beta^{-(k-h)}$  and every cylinder with digits  $\epsilon_0, \dots, \epsilon_{k-h+1}$  has measure  $\beta^{-(k-h+2)}$ . Thus it follows

$$\mu(Z) = \beta^{-(k-h+1)} - (a-1)\beta^{-(k-h+2)}.$$

Considering  $\phi_\beta(Z)$  we have

$$\phi_\beta(Z) = \left[ \sum_{i=0}^{k-h+1} \frac{\epsilon_i}{\beta^{i+1}}, \sum_{i=0}^{k-h} \frac{\epsilon_i}{\beta^{i+1}} + \frac{(\epsilon_{k-h-1} + 1)}{\beta^{k-h+2}} \right),$$

and thus  $\lambda(\phi_\beta(Z)) = \mu(Z)$ .  $\square$

As mentioned in the previous section, a result of Frougny and Solomyak [12, Lemma 3] implies that the dominant root of

$$x^2 - a_0x - a_1, \quad a_0, a_1 \geq 1,$$

is a Pisot number if and only if  $a_0 \geq a_1$ . Lemma 1 shows that the image of  $\mathcal{K}_G^0$  under  $\phi_\beta$  is not a subset of  $[0, 1)$ , when  $a_0 > a_1$ . Thus Lemma 2 characterizes all van der Corput-type constructions for  $d = 2$ .

**Theorem 3.** *Let  $G^1, \dots, G^s$  be numeration systems as in Theorem 2 and let  $\beta_1, \dots, \beta_s$  denote the roots of the corresponding characteristic equations. Then the  $s$ -dimensional  $\beta$ -adic Halton sequence  $(\phi_\beta(n))_{n \in \mathbb{N}}$  is u.d. in  $[0, 1)^s$ .*

*Proof.* By Lemma 2 and the definition of the Monna map we obtain an isometry between the dynamical systems  $((\mathcal{K}_{G^1}, \tau_1) \times \dots \times (\mathcal{K}_{G^s}, \tau_s))$  and  $(([0, 1], T_1) \times \dots \times ([0, 1], T_s))$  where

$$T_i: [0, 1] \rightarrow [0, 1], \quad T_i(x) := \phi_{\beta_i} \circ \tau_i \circ \phi_{\beta_i}^{-1}(x).$$

Let  $\mathbf{T}\mathbf{x} = (T_1x_1, \dots, T_sx_s)$  for  $\mathbf{x} = (x_1, \dots, x_s) \in [0, 1]^s$ . Hence by Birkhoff's ergodic theorem,  $(\mathbf{T}^n\mathbf{x})_{n \in \mathbb{N}}$  is u.d. in  $[0, 1]^s$  for all  $\mathbf{x} \in [0, 1]^s$ . In particular,  $(\phi_{\beta}(n))_{n \in \mathbb{N}} = (\mathbf{T}^n\mathbf{0})_{n \in \mathbb{N}}$  is u.d.  $\square$

Note that the classical  $\mathbf{b}$ -adic Halton sequence with pairwise coprime integer bases  $b_1, \dots, b_s \geq 2$ , is covered by Theorem 3.

**Theorem 4.** *Let the numeration system  $G$  be defined by the coefficients  $(a_0, a_1, a_2) = (1, 0, 1)$ ,  $\beta$  be its characteristic root and  $\tau$  the odometer on  $G$ . Then  $\mu(Z) = \lambda(\phi_{\beta}(Z))$  for all cylinder sets  $Z$ . Thus  $T(x) = \phi_{\beta} \circ \tau \circ \phi_{\beta}^{-1}(x)$  is uniquely ergodic and  $(T^n x)_{n \in \mathbb{N}}$  is u.d. for all  $x$  in  $[0, 1]$ . Furthermore the spectrum of  $T$  is given by*

$$\Gamma = \left\{ \exp\left(2\pi i \frac{c}{\beta^l}\right) : m, l, c \in \mathbb{N} \cup \{0\} \right\}. \quad (19)$$

*Proof.* It is well known that  $\beta$  is a Pisot number and equation (7) holds since  $[\beta] = 1 = a_0$ . Hypothesis 1 was proved for this case in [6, Theorem 4]. The proof that Hypothesis 2 is fulfilled can be found in [2, Theorem 3]. Equation (19) follows from the proof of Theorem 2.

Now we have to prove that  $\phi_{\beta}$  transports the measure  $\mu$  to the Lebesgue measure on  $[0, 1]$ . First we assume  $k \geq 3$ . Let the cylinder  $Z$  be defined by the fixed digits  $\epsilon_0, \dots, \epsilon_{k-1}$ . We consider four different cases, first  $\epsilon_{k-3} = \epsilon_{k-2} = \epsilon_{k-1} = 0$ . Then  $F_{k,0} = 1, F_{k,1} = 2, F_{k,2} = 3$  and we get by (9) that

$$\mu(Z) = \beta^{-k}.$$

Furthermore, by the same argument as in the first part of the proof of Theorem 3 we obtain

$$\phi_{\beta}(Z) = \left[ \sum_{i=0}^{k-1} \frac{\epsilon_i}{\beta^{i+1}}, \sum_{i=0}^{k-2} \frac{\epsilon_i}{\beta^{i+1}} + \frac{(\epsilon_{k-1} + 1)}{\beta^k} \right),$$

and thus  $\lambda(\phi_{\beta}(Z)) = \beta^{-k}$ .

Now let  $\epsilon_{k-3} = 1, \epsilon_{k-2} = \epsilon_{k-1} = 0$ . Hence  $F_{k,0} = 1, F_{k,1} = 2, F_{k,2} = 3$  and  $\mu(Z) = \beta^{-k}$ . We have

$$\begin{aligned} \phi_{\beta}(Z) &= \left[ \sum_{i=0}^{k-1} \frac{\epsilon_i}{\beta^{i+1}}, \sum_{i=0}^{k-1} \frac{\epsilon_i}{\beta^{i+1}} + \beta^{-k} \sum_{i=0}^{\infty} \beta^{-(3i+1)} \right) \\ &= \left[ \sum_{i=0}^{k-1} \frac{\epsilon_i}{\beta^{i+1}}, \sum_{i=0}^{k-1} \frac{\epsilon_i}{\beta^{i+1}} + \beta^{-k} \right), \end{aligned}$$

thus again  $\lambda(\phi_{\beta}(Z)) = \beta^{-k}$ . Now assume  $\epsilon_{k-2} = 1, \epsilon_{k-1} = 0$ . Hence  $F_{k,0} = 1, F_{k,1} = 1, F_{k,2} = 2$  and

$$\mu(Z) = \beta^{-k} \frac{\beta^{-2} + 1}{\beta^{-2} + \beta^{-1} + 1}.$$

Similarly as above we obtain

$$\begin{aligned}\phi_\beta(Z) &= \left( \sum_{i=0}^{k-1} \frac{\epsilon_i}{\beta^{i+1}}, \sum_{i=0}^{k-1} \frac{\epsilon_i}{\beta^{i+1}} + \beta^{-(k+1)} \sum_{i=0}^{\infty} \beta^{-(3i+1)} \right) \\ &= \left( \sum_{i=0}^{k-1} \frac{\epsilon_i}{\beta^{i+1}}, \sum_{i=0}^{k-1} \frac{\epsilon_i}{\beta^{i+1}} + \beta^{-(k+1)} \right),\end{aligned}$$

thus  $\lambda(\phi_\beta(Z)) = \beta^{-(k+1)}$ . Now we have

$$\beta^{-(k+1)} = \beta^{-k} \frac{\beta^{-2} + 1}{\beta^{-2} + \beta^{-1} + 1}$$

which is equivalent to  $\beta^{-3} + \beta^{-2} + \beta^{-1} = \beta^{-2} + 1$  and holds by (5). In the last case we assume  $\epsilon_{k-1} = 1$ , thus  $F_{k,0} = F_{k,1} = F_{k,2} = 1$  and

$$\mu(Z) = \beta^{-k} \frac{1}{\beta^{-2} + \beta^{-1} + 1}.$$

As above we get  $\lambda(\phi_\beta(Z)) = \beta^{-(k+2)}$  and the result follows since

$$\beta^{-(k+2)} = \beta^{-k} \frac{1}{\beta^{-2} + \beta^{-1} + 1}$$

is equivalent to  $\beta^{-3} + \beta^{-1} = 1$ . The cases, where  $k < 3$ , follow by the same arguments.  $\square$

As a consequence of Theorem 1 we can construct uniformly distributed two-dimensional sequences  $(\phi_{\beta_1}(n), \phi_{\beta_2}(n))_{n \in \mathbb{N}}$ , where  $\beta_1$  is the characteristic root in Theorem 4,  $\beta_2$  is the characteristic root of a numeration system in Theorem 2 and  $\frac{\beta_1^k}{\beta_2^l} \notin \mathbb{Q}$  for all integers  $k, l > 0$ .

Theorem 4 extends the examples given in [5, Proposition 13,14], where the authors consider  $G$ -additive functions which lead to u.d. point sequences in the unit interval. Furthermore, it is possible to show that the one-dimensional point sequence in the previous theorem is a low-discrepancy sequence by mimicking the proof for the  $b$ -adic van der Corput sequence, see e.g. [20, 5, 7].

In [8], the authors present the so-called Kakutani-Fibonacci transformation, proving ergodicity on the unit interval and showing that the orbit of 0 is exactly the LS-sequence with parameters  $L = S = 1$  defined in [7]. With our different approach we can show that this transformation is uniquely ergodic, thus the orbit of  $x$  under the transformation is u.d. for every  $x \in [0, 1)$ .


The Kakutani-Fibonacci transformation was first introduced in [13] and discussed in detail in [8] and [13]. In the following we give an informal description.

Let us consider the unit interval  $[0, 1)$  and split it into two consecutive subintervals  $[0, \alpha)$  and  $[\alpha, 1)$  of lengths  $\alpha$  and  $\alpha^2$ , respectively, where  $\alpha$  is the reciprocal golden ratio  $\frac{\sqrt{5}-1}{2}$ . So we obtain two systems of half-open intervals, in the first step  $L_1 = \{[0, \alpha)\}$  (“long”) and  $S_1 = \{[\alpha, 1)\}$  (“short”). One can easily iterate

this procedure in the following way. Let us split all intervals contained in  $L_1$ , proportionally to  $\alpha^2$  and  $\alpha^3$ , respectively. Thus we obtain two systems of half-open intervals  $L'_2$  and  $S'_2$ , the first one containing intervals of length  $\alpha^2$  and the second of length  $\alpha^3$ , respectively.

This yields two new systems of half-open intervals,  $L_2 = \{L'_2 \cup S_1\}$ ,  $S_2 = S'_2$ . Note that the ordering of the intervals is essential and follows from the illustration in the picture below.

In general, at the  $k$ -th step, we split only the intervals in  $L_{k-1}$ , proportionally to  $\alpha^k$  and  $\alpha^{k+1}$  forming two systems of half-open intervals  $L'_k$  and  $S'_k$ , the first one containing intervals of length  $\alpha^k$  and the second of length  $\alpha^{k+1}$ , respectively. Again this yields a system of half-open intervals  $L_k = \{L'_k \cup S_{k-1}\}$  and  $S_k = S'_k$ .



This construction procedure is well-known as cutting-stacking method and the transformation associated to it is defined as the translation from the first interval of  $L_k$  onto its second one, the second onto the third and so on till the last interval of the column  $L_k$ . Then the transformation is given as a contraction with factor  $\beta$  from the top of  $L_k$  to the bottom of  $S_k$ , and then again as the translation from the first interval of  $S_k$  onto the same and so on and so forth.

In the sequel we give an explicit expression for the Kakutani-Fibonacci transformation  $T$ , which is the limit transformation of the above construction, see [8].

**Definition 2.** *The Kakutani-Fibonacci transformation is a piecewise linear map  $T : [0, 1) \rightarrow [0, 1)$ , whose restriction to  $I_k$  is  $T_k$ , where*

$$T_1 x = x + \alpha \quad \text{if } x \in I_1 = [0, \alpha^2)$$

and, for every  $k \geq 1$ ,

$$T_{2k} x = x + \alpha^{2k} - \sum_{j=0}^{k-1} \alpha^{2j+1} \quad \text{if } x \in I_{2k} = \left[ \sum_{j=0}^{k-1} \alpha^{2j+1}, \sum_{j=0}^k \alpha^{2j+1} \right)$$

and

$$T_{2k+1} x = x + \alpha^{2k+1} - \sum_{j=0}^{k-1} \alpha^{2(j+1)} \quad \text{if } x \in I_{2k+1} = \left[ \sum_{j=0}^{k-1} \alpha^{2(j+1)}, \sum_{j=0}^k \alpha^{2(j+1)} \right).$$

For proving unique ergodicity of the Kakutani-Fibonacci transformation we need the following lemma.

**Lemma 3.** *Let  $\beta$  be the golden ratio  $\frac{\sqrt{5}+1}{2}$ . Then we have  $Tx = \phi_\beta \circ \tau \circ \phi_\beta^{-1}x$  for all  $\beta$ -adic rationals*

$$x = \sum_{i=1}^k \frac{\epsilon_i}{\beta^i}$$

with coefficients  $\epsilon_i \in \{0, 1\}$ .

*Proof.* This follows from [1, Lemma 3] specializing  $L = S = 1$ . Applying this lemma combined with Equation 2 in Section 2 of [1] yields that the  $\beta$ -adic Monna is bijective on the set of all  $\beta$ -adic rationals. Combining this with [8, Theorem 16] completes the proof of Lemma 3.  $\square$

**Theorem 5.** *The Kakutani-Fibonacci transformation is uniquely ergodic.*

*Proof.* By Theorem 3,  $\phi_\beta \circ \tau \circ \phi_\beta^{-1}$  is uniquely ergodic, thus by Lemma 3 and the denseness of the  $\beta$ -adic rationals  $T$  is uniquely ergodic.  $\square$

## Acknowledgements

We are grateful to P. Grabner, P. Liardet and the anonymous referee for various helpful remarks and comments.

## References

- [1] C. Aistleitner, M. Hofer, and V. Ziegler. On the uniform distribution modulo 1 of multidimensional LS-sequences. *to appear: Annali di Matematica Pura ed Applicata*, 2013.

- [2] S. Akiyama. Cubic Pisot numbers with finite beta expansions. *In F. Halter-Koch and R. Tichy (eds.): Algebraic Number Theory and Diophantine Analysis*, pages 11–26, 2000.
- [3] S. Akiyama, H. Brunotte, A. Pethö, and J. M. Thuswaldner. Generalized radix representations and dynamical systems II. *Acta Arithmetica*, 121:21–61, 2006.
- [4] S. Asmussen and H. Albrecher. *Ruin Probabilities, 2nd eds.* World Scientific, Singapore, 2010.
- [5] G. Barat and P. Grabner. Distribution properties of G-additive functions. *Journal of Number Theory*, 60:103–123, 1996.
- [6] H. Bruin, G. Keller, and M. Pierre. Adding machines and wild attractors. *Ergodic Theory and Dynamical Systems*, 17(6):1267–1287, 1997.
- [7] I. Carbone. Discrepancy of LS-sequences of partitions and points. *Ann. Mat. Pura Appl. (4)*, 191(4):819–844, 2012.
- [8] I. Carbone, M. R. Iacò, and A. Volčič. A dynamical system approach to the Kakutani-Fibonacci sequence. *to appear: Ergodic Theory and Dynamical Systems*, 2012.
- [9] M. Drmota and M. Infusino. On the discrepancy of some generalized Kakutani’s sequences of partitions. *Uniform Distribution Theory*, 7:75–104, 2012.
- [10] M. Drmota and R. F. Tichy. *Sequences, discrepancies and applications*, volume 1651 of *Lecture Notes in Mathematics*. Springer-Verlag, Berlin, 1997.
- [11] A. S. Fraenkel. Systems of numeration. *Amer. Math. Monthly*, 92(2):105–114, 1985.
- [12] C. Frougny and B. Solomyak. Finite beta-expansions. *Ergodic Theory and Dynamical Systems*, 12:713–723, 1992.
- [13] P. Grabner, P. Hellekalek, and P. Liardet. The dynamical point of view of low-discrepancy sequences. *Uniform Distribution Theory*, 7(1):11–70, 2012.
- [14] P. Grabner, P. Liardet, and R. Tichy. Odometers and systems of numeration. *Acta Arithmetica*, 70:103–123, 1995.
- [15] P. Grabner and R. Tichy. Contributions to Digit Expansions with Respect to Linear Recurring Sequences. *Journal of Number Theory*, 36:160–169, 1990.
- [16] P. Grabner and R. Tichy.  $\alpha$ -expansions, linear recurrences, and the sum-of-digits function. *Manuscripta Mathematica*, 70:311–324, 1991.
- [17] J. H. Halton. On the efficiency of certain quasi-random sequences of points in evaluating multi-dimensional integrals. *Numer. Math.*, 2:84–90, 1960.
- [18] E. Hlawka. Funktionen von beschränkter Variation in der Theorie der Gleichverteilung. *Annali di Matematica Pura ed Applicata*, (54):325–333, 1961.

- [19] M. Hollander. *Linear numeration systems, finite beta-expansions, and discrete spectrum of substitution dynamical systems*. PhD Thesis, Washington University, 1996.
- [20] L. Kuipers and H. Niederreiter. *Uniform distribution of sequences*. Wiley-Interscience [John Wiley & Sons], New York, 1974.
- [21] M. Mori and M. Mori. Dynamical systems generated by algebraic method and low discrepancy sequences. *Monte Carlo Methods and Applications*, 18:327–351, 2012.
- [22] S. Ninomiya. Constructing a new class of low-discrepancy sequences by using the  $\beta$ -adic transformation. *Math. Comput. Simulation*, 47(2):403–418, 1998.
- [23] G. Ökten, B. Tuffin, and V. Burago. A central limit theorem and improved error bounds for a hybrid-Monte Carlo sequence with applications in computational finance. *Journal of Complexity*, 22(4):435–458, 2006.
- [24] W. Parry. On the  $\beta$ -expansions of real numbers. *Acta Math. Acad. Sci. Hungar.*, 11:401–416, 1960.
- [25] B. Solomyak. Substitutions, adic transformations and beta-expansions. *in: Contemporary Mathematics (Proc. of the Conference on Symbolic Dynamics, Yale, 1991)*, 135:361–372, 1992.
- [26] W. Steiner. Regularities of the distribution of  $\beta$ -adic van der Corput sequences. *Monatshefte Mathematik*, 149(1):67–81, 2006.
- [27] W. Steiner. Regularities of the distribution of abstract van der Corput sequences. *Uniform Distribution Theory*, 4(2):81–100, 2009.
- [28] P. Walters. *An introduction to ergodic theory*, volume 79 of *Graduate Texts in Mathematics*. Springer-Verlag, New York, 1982.
- [29] H. Weyl. Über die Gleichverteilung von Zahlen mod. Eins. *Mathematische Annalen*, 77:313–352, 1916.