

# Selfadjoint Schrödinger operators on the half-space with compactly supported Robin boundary conditions

Jussi Behrndt<sup>1</sup>, Christian Kühn<sup>2,\*</sup>, and Jonathan Rohleder<sup>1</sup>

<sup>1</sup> Institut für Numerische Mathematik, Technische Universität Graz, Steyrergasse 30, 8010 Graz, Austria

<sup>2</sup> Institut für Mathematik, MA 6-4, Technische Universität Berlin, Straße des 17. Juni 136, 10623 Berlin, Germany

Schrödinger differential operators on a half-space with compactly supported Robin boundary conditions are studied.

Copyright line will be provided by the publisher

## 1 Introduction

We investigate realizations of the differential expression  $-\Delta + V$  on the half-space  $\mathbb{R}_+^n = \{(x', x_n) : x' \in \mathbb{R}^{n-1}, x_n > 0\}$ ,  $n \geq 2$ , with a real-valued, bounded potential  $V$ . More precisely, we study the differential operator

$$A_g u = -\Delta u + V u, \quad \text{dom } A_g = \left\{ u \in H_{\Delta}^{3/2}(\mathbb{R}_+^n) : \partial_{\nu} u|_{\mathbb{R}^{n-1}} = g \cdot (u|_{\mathbb{R}^{n-1}}) \right\}, \quad (1)$$

in  $L^2(\mathbb{R}_+^n)$ , where  $H_{\Delta}^{3/2}(\mathbb{R}_+^n) = \{u \in H^{3/2}(\mathbb{R}_+^n) : \Delta u \in L^2(\mathbb{R}_+^n)\}$  and  $g : \mathbb{R}^{n-1} \rightarrow \mathbb{R}$  is a bounded, real function with compact support. The aim of the present note is to show that  $A_g$  is a selfadjoint, compact perturbation in the resolvent sense of the selfadjoint realization  $A_N$  of  $-\Delta + V$  with Neumann boundary conditions. In particular this guarantees that  $A_g$  and  $A_N$  have the same essential spectrum. We point out that the latter can still be proved under slightly weaker assumptions on  $g$ , see [10] for a more general approach and [8] for a result with a more regular  $g$  in dimension  $n = 2$ . Our proofs make use of techniques which were originally developed in [2, 3] for the treatment of elliptic differential operators on domains with a compact boundary. For further recent developments in this area we refer the reader to [1, 4, 6, 11, 12].

## 2 Preliminaries

In this section we fix some notation and recall some known facts on Sobolev spaces and Schrödinger operators; proofs and further details can be found in [9] and, e.g., [7, Chapter 9]. Let  $K \subset \mathbb{R}^{n-1}$  be a compact set and let  $H^s(\mathbb{R}_+^n)$  and  $H^s(K) = \{f|_K : f \in H^s(\mathbb{R}^{n-1})\}$  be the Sobolev spaces of order  $s > 0$  on  $\mathbb{R}_+^n$  and  $K$ , respectively. For  $u \in H_{\Delta}^{3/2}(\mathbb{R}_+^n)$  we denote by  $u|_{\mathbb{R}^{n-1}}$  the trace of  $u$  on the boundary  $\mathbb{R}^{n-1}$  of  $\mathbb{R}_+^n$ , by  $\partial_{\nu} u|_{\mathbb{R}^{n-1}} = -\frac{\partial u}{\partial x_n}|_{\mathbb{R}^{n-1}}$  the derivative of  $u$  along the outer normal vector field on  $\mathbb{R}^{n-1}$ , and by  $u|_K$  and  $\partial_{\nu} u|_K$ ,  $\partial_{\nu} u|_{\mathbb{R}^{n-1} \setminus K}$  their restrictions to  $K$  and  $\mathbb{R}^{n-1} \setminus K$ , respectively. The mappings  $\Gamma_0$  and  $\Gamma_1$  given by

$$\Gamma_0 : H_{\Delta}^{3/2}(\mathbb{R}_+^n) \rightarrow L^2(K), \quad \Gamma_0 u = \partial_{\nu} u|_K \quad \text{and} \quad \Gamma_1 : H_{\Delta}^{3/2}(\mathbb{R}_+^n) \rightarrow H^1(K), \quad \Gamma_1 u = u|_K \quad (2)$$

are surjective.

Here and in the following let  $V \in L^{\infty}(\mathbb{R}_+^n)$  be real-valued. It is well known that the *Neumann operator*

$$A_N u = -\Delta u + V u, \quad \text{dom } A_N = \left\{ u \in H_{\Delta}^{3/2}(\mathbb{R}_+^n) : \partial_{\nu} u|_{\mathbb{R}^{n-1}} = 0 \right\}$$

is a selfadjoint realization of  $-\Delta + V$  in  $L^2(\mathbb{R}_+^n)$ , and by elliptic regularity  $\text{dom } A_N \subset H^2(\mathbb{R}_+^n)$  holds. Note that this yields the decomposition  $\{u \in H_{\Delta}^{3/2}(\mathbb{R}_+^n) : \partial_{\nu} u|_{\mathbb{R}^{n-1} \setminus K} = 0\} = \text{dom } A_N \dot{+} \mathcal{N}_{\lambda}$  for each  $\lambda$  in the resolvent set  $\rho(A_N)$  of  $A_N$ , where  $\mathcal{N}_{\lambda} := \{u \in H^{3/2}(\mathbb{R}_+^n) : -\Delta u + V u = \lambda u, \partial_{\nu} u|_{\mathbb{R}^{n-1} \setminus K} = 0\}$ . This, together with (2), ensures that the *Poisson operator*

$$\gamma(\lambda) : L^2(K) \rightarrow L^2(\mathbb{R}_+^n), \quad \partial_{\nu} u_{\lambda}|_K \mapsto u_{\lambda}, \quad u_{\lambda} \in \mathcal{N}_{\lambda}, \quad (3)$$

and the *Neumann-to-Dirichlet operator*

$$M(\lambda) : L^2(K) \rightarrow L^2(K), \quad \partial_{\nu} u_{\lambda}|_K \mapsto u_{\lambda}|_K, \quad u_{\lambda} \in \mathcal{N}_{\lambda}, \quad (4)$$

are well-defined for each  $\lambda \in \rho(A_N)$ . Moreover,  $\gamma(\lambda)$  and  $M(\lambda)$  are bounded and  $\text{ran } M(\lambda) = H^1(K)$  holds.

\* Corresponding author: Email ckuehn@mail.math.tu-berlin.de

### 3 Selfadjoint Schrödinger operators on the half-space

The following theorem is the main result of this note. For  $g \in L^\infty(\mathbb{R}^{n-1})$ ,  $\text{supp } g = K$ , we denote by  $G$  the operator of multiplication with the function  $g|_K$  in  $L^2(K)$ .

**Theorem 3.1** *Let  $K \subset \mathbb{R}^{n-1}$  be a compact set and let  $g \in L^\infty(\mathbb{R}^{n-1})$  be a real-valued function with  $\text{supp } g = K$ . Then the operator  $A_g$  in (1) is selfadjoint in  $L^2(\mathbb{R}_+^n)$  and  $\lambda \in \rho(A_N)$  is an eigenvalue of  $A_g$  if and only if 1 is an eigenvalue of  $GM(\lambda)$ . The resolvent difference*

$$(A_g - \lambda)^{-1} - (A_N - \lambda)^{-1} = \gamma(\lambda)(I - GM(\lambda))^{-1}G\gamma(\bar{\lambda})^*, \quad \lambda \in \rho(A_g) \cap \rho(A_N), \quad (5)$$

*is compact and, in particular, the essential spectra of  $A_g$  and  $A_N$  coincide.*

**Proof.** Let us first show that  $\lambda \in \rho(A_N)$  is an eigenvalue of  $A_g$  if and only if 1 is an eigenvalue of  $GM(\lambda)$ . For  $u \in \ker(A_g - \lambda)$ ,  $u \neq 0$ , we have  $\Gamma_0 u \neq 0$  and  $GM(\lambda)\Gamma_0 u = G\Gamma_1 u = \Gamma_0 u$ . Thus  $I - GM(\lambda)$  is not injective. Conversely,  $f \in \ker(I - GM(\lambda))$ ,  $f \neq 0$ , implies  $\gamma(\lambda)f \in \text{dom } A_g$ ,  $(A_g - \lambda)\gamma(\lambda)f = 0$ , and  $\gamma(\lambda)f \neq 0$ . Thus  $\gamma(\lambda)f$  is an eigenfunction of  $A_g$  corresponding to the eigenvalue  $\lambda$ .

Next we show that  $A_g$  is a selfadjoint operator in  $L^2(\mathbb{R}_+^n)$ . For this note first that for  $u \in \text{dom } A_g$  we have

$$(A_g u, u) = \int_{\mathbb{R}_+^n} (-\Delta + V)u \bar{u} dx = \int_{\mathbb{R}_+^n} |\nabla u|^2 + V|u|^2 dx - \int_K g|u|^2 d\sigma \in \mathbb{R},$$

so that  $A_g$  is a symmetric in operator in  $L^2(\mathbb{R}_+^n)$ . Hence it is sufficient to verify that  $A_g - \lambda$  is surjective for  $\lambda \in \mathbb{C} \setminus \mathbb{R}$ . Fix some  $\lambda \in \mathbb{C} \setminus \mathbb{R}$ , choose an arbitrary  $u \in L^2(\mathbb{R}_+^n)$ , and define

$$v := (A_N - \lambda)^{-1}u + \gamma(\lambda)(I - GM(\lambda))^{-1}G\gamma(\bar{\lambda})^*u. \quad (6)$$

In the following we will show that  $v$  is well-defined and belongs to  $\text{dom } A_g$  with  $(A_g - \lambda)v = u$ . The operator  $\gamma(\lambda)$  and hence also  $\gamma(\bar{\lambda})^*$  and  $G\gamma(\bar{\lambda})^*$  are bounded and everywhere defined. Furthermore, since  $\text{ran } M(\lambda) = H^1(K)$  and the embedding from  $H^1(K)$  into  $L^2(K)$  is compact,  $M(\lambda)$  and  $GM(\lambda)$  are also compact operators in  $L^2(K)$ . Together with the fact that 1 is not an eigenvalue of  $GM(\lambda)$  we conclude that the operator  $I - GM(\lambda)$  has an everywhere defined, bounded inverse, i.e.,  $v$  in (6) is well-defined. From the definition of  $v$  it is easy to see that  $v \in H_{\Delta}^{3/2}(\mathbb{R}_+^n)$  and  $\partial_\nu v|_{\mathbb{R}^{n-1} \setminus K} = 0$  holds.

It remains to show  $G\Gamma_1 v = \Gamma_0 v$  and  $(A_g - \lambda)v = u$ . In fact, as a consequence of the second Green identity we find  $\Gamma_1(A_N - \lambda)^{-1}u = \gamma(\bar{\lambda})^*u$  and therefore we conclude from (6)

$$G\Gamma_1 v = G\gamma(\bar{\lambda})^*u + GM(\lambda)(I - GM(\lambda))^{-1}G\gamma(\bar{\lambda})^*u = (I - GM(\lambda))^{-1}G\gamma(\bar{\lambda})^*u = \Gamma_0 v.$$

Thus we have shown  $v \in \text{dom } A_g$  and from  $(A_g - \lambda)v = (-\Delta + V - \lambda)v = u$  we obtain that  $A_g - \lambda$  is surjective and, hence,  $A_g$  is selfadjoint. Moreover, we have shown the formula (5) for all  $\lambda \in \mathbb{C} \setminus \mathbb{R}$  and the same reasoning applies for real  $\lambda \in \rho(A_N) \cap \rho(A_g)$ . As mentioned above,  $\gamma(\bar{\lambda})^* = \Gamma_1(A_N - \lambda)^{-1}$ , in particular,  $\text{ran } \gamma(\bar{\lambda})^* \subset H^{3/2}(K)$ , which is compactly embedded in  $L^2(K)$ . This shows that the right hand side in (5) is compact for  $\lambda \in \mathbb{C} \setminus \mathbb{R}$ . Hence  $(A_g - \lambda)^{-1} - (A_N - \lambda)^{-1}$  is compact for each  $\lambda \in \rho(A_g) \cap \rho(A_N)$ , and, in particular,  $A_g$  and  $A_N$  have the same essential spectrum.  $\square$

We obtain the following corollary in the case  $V = 0$ .

**Corollary 3.2** *Let  $V = 0$ . Then the essential spectrum of the operator  $A_g$  in (1) is given by  $[0, +\infty)$ . Moreover,  $\lambda < 0$  is an eigenvalue of  $A_g$  if and only if 1 is an eigenvalue of  $G\iota^*(-\Delta_{\mathbb{R}^{n-1}} - \lambda)^{-1/2}\iota$ , where  $\iota$  denotes the embedding from  $L^2(K)$  into  $L^2(\mathbb{R}^{n-1})$  and  $\Delta_{\mathbb{R}^{n-1}}$  is the Laplacian on  $\mathbb{R}^{n-1}$ .*

**Proof.** In the case  $V = 0$  it is well-known that the spectrum and essential spectrum of  $A_N$  is given by  $[0, +\infty)$ . Moreover, one computes similarly as in [7, Chapter 9] that  $M(\lambda) = \iota^*(-\Delta_{\mathbb{R}^{n-1}} - \lambda)^{-1/2}\iota$  holds.  $\square$

### References

- [1] H. Abels, G. Grubb, and I. G. Wood, submitted.
- [2] J. Behrndt and M. Langer, *J. Funct. Anal.* **243** (2007), 536–565.
- [3] J. Behrndt, M. Langer, and V. Lotoreichik, submitted.
- [4] B. M. Brown, G. Grubb, and I. G. Wood, *Math. Nachr.* **282** (2009), 314–347.
- [5] F. Gesztesy and M. Mitrea, *J. Differential Equations* **247** (2009), 2871–2896.
- [6] F. Gesztesy and M. Mitrea, *J. Anal. Math.* **113** (2011), 53–172.
- [7] G. Grubb, *Distributions and Operators* (Springer, New York, 2009).
- [8] M. Jířek, *SIGMA Symmetry Integrability Geom. Methods Appl.* **3** (2007), 12 pp.
- [9] J. L. Lions and E. Magenes, *Non-Homogeneous Boundary Value Problems and Applications I* (Springer, Berlin, 1972).
- [10] V. Lotoreichik and J. Rohleder, submitted.
- [11] M. M. Malamud, *Russ. J. Math. Phys.* **17** (2010), 96–125.
- [12] A. Posilicano and L. Raimondi, *J. Phys. A: Math. Theor.* **42** (2009), 11 pp.