

FINITE ROGERS-RAMANUJAN TYPE CONTINUED FRACTIONS

HELMUT PRODINGER

Dedicated to Peter Paule on the occasion of his 60th birthday

ABSTRACT. New finite continued fractions related to Bressoud and Santos polynomials are established.

1. INTRODUCTION

Define, as it is common today, $(x; q)_n := (1-x)(1-xq)\dots(1-xq^{n-1})$, where we assume that $|q| < 1$, and we allow n also to be 0 and infinity. We also need the coefficients $\begin{bmatrix} n \\ k \end{bmatrix} := \frac{(q; q)_n}{(q; q)_k (q; q)_{n-k}}$. These standard notations can be found e. g. in the classic book [1].

The two Rogers-Ramanujan identities [6, 1]

$$\sum_{n \geq 0} \frac{q^{n^2}}{(q; q)_n} = \frac{1}{(q; q^5)_\infty (q^4; q^5)_\infty},$$

$$\sum_{n \geq 0} \frac{q^{n^2+n}}{(q; q)_n} = \frac{1}{(q^2; q^5)_\infty (q^3; q^5)_\infty}$$

are very popular, influential, useful and historically interesting. Let

$$F(z) = \sum_{n \geq 0} \frac{q^{n^2} z^n}{(q; q)_n} \quad \text{and} \quad G(z) = \sum_{n \geq 0} \frac{q^{n^2+n} z^n}{(q; q)_n};$$

the continued fraction (due to Ramanujan)

$$\frac{zG(z)}{F(z)} = \frac{z}{1 + \frac{zq}{1 + \frac{zq^2}{1 + \dots}}}$$

is also very well known, see [5, Entry 5] and [6, (6.1)].

There are several families of polynomials that approximate $F(z)$ and $G(z)$. Probably the the most well known are

$$f_n(z) = \sum_{j \geq 0} q^{j^2} \begin{bmatrix} n+1-j \\ j \end{bmatrix} z^j \rightarrow F(z) \quad \text{and} \quad g_n(z) = \sum_{j \geq 0} q^{j^2+j} \begin{bmatrix} n-j \\ j \end{bmatrix} z^j \rightarrow G(z) \quad \text{for } n \rightarrow \infty,$$

Key words and phrases. Bressoud polynomials, Santos polynomials, Rogers-Ramanujan identities.

because of their link to the Schur polynomials, see [1].

The finite continued fraction

$$\frac{z g_n(z)}{f_n(z)} = \frac{z}{1 + \frac{zq}{1 + \frac{zq^2}{\ddots 1 + \frac{zq^n}{1}}}}$$

is also known [5, Entry 16].

The polynomials

$$s_n(z) = \sum_{j \geq 0} q^{j^2} \begin{bmatrix} n \\ j \end{bmatrix} z^j \rightarrow F(z) \quad \text{and} \quad t_n(z) = \sum_{j \geq 0} q^{j^2+j} \begin{bmatrix} n \\ j \end{bmatrix} z^j \rightarrow G(z) \quad \text{for } n \rightarrow \infty,$$

due to *Bressoud* [7], see also [8] are less well known.

2. BRESSOUD POLYNOMIALS AND CONTINUED FRACTIONS

We will establish the following attractive finite continued fraction:

$$\frac{z t_n(z)}{s_n(z)} = \frac{z}{1 + \frac{zq(1-q^n)}{1 + \frac{zq^2}{1 + \frac{zq^3(1-q^{n-1})}{1 + \frac{zq^4}{1 + \ddots}}}}}$$

To prove this statement by induction, define the righthand side by $T_n(z)$. It is plain to see that $T_0(z) = z$, and

$$T_n(z) = \frac{z}{1 + \frac{zq(1-q^n)}{1 + T_{n-1}(zq^2)}}.$$

We are left to prove that

$$\begin{aligned} \frac{z t_n(z)}{s_n(z)} &= \frac{z}{1 + \frac{zq(1-q^n)}{1 + \frac{z t_{n-1}(zq^2)}{s_{n-1}(zq^2)}}} = \frac{z}{1 + \frac{zq(1-q^n)s_{n-1}(zq^2)}{s_{n-1}(zq^2) + z t_{n-1}(zq^2)}} \\ &= \frac{z(s_{n-1}(zq^2) + z t_{n-1}(zq^2))}{s_{n-1}(zq^2) + z t_{n-1}(zq^2) + zq(1-q^n)s_{n-1}(zq^2)} \end{aligned}$$

which amounts to prove that

$$\begin{aligned} t_n(z) &= s_{n-1}(zq^2) + zt_{n-1}(zq^2), \\ s_n(z) &= s_{n-1}(zq^2) + zt_{n-1}(zq^2) + zq(1-q^n)s_{n-1}(zq^2). \end{aligned}$$

We will show that the coefficients of z^j coincide, which is trivial for $j = 0$, so we assume $j \geq 1$:

$$q^{j^2+j} \begin{bmatrix} n \\ j \end{bmatrix} = q^{j^2+2j} \begin{bmatrix} n-1 \\ j \end{bmatrix} + q^{(j-1)^2+3(j-1)+2} \begin{bmatrix} n-1 \\ j-1 \end{bmatrix},$$

which is equivalent to

$$\begin{bmatrix} n \\ j \end{bmatrix} = q^j \begin{bmatrix} n-1 \\ j \end{bmatrix} + \begin{bmatrix} n-1 \\ j-1 \end{bmatrix}$$

and therefore true. The second one goes like this:

$$q^{j^2} \begin{bmatrix} n \\ j \end{bmatrix} = q^{j^2+2j} \begin{bmatrix} n-1 \\ j \end{bmatrix} + q^{(j-1)^2+3(j-1)+2} \begin{bmatrix} n-1 \\ j-1 \end{bmatrix} + q(1-q^n)q^{(j-1)^2+2(j-1)} \begin{bmatrix} n-1 \\ j-1 \end{bmatrix},$$

which is equivalent to

$$\begin{bmatrix} n \\ j \end{bmatrix} = q^{2j} \begin{bmatrix} n-1 \\ j \end{bmatrix} + q^j \begin{bmatrix} n-1 \\ j-1 \end{bmatrix} + (1-q^j) \begin{bmatrix} n \\ j \end{bmatrix},$$

and further to

$$\begin{bmatrix} n \\ j \end{bmatrix} = q^j \begin{bmatrix} n-1 \\ j \end{bmatrix} + \begin{bmatrix} n-1 \\ j-1 \end{bmatrix},$$

which finishes the proof.

3. IDENTITIES 39 AND 38 FROM SLATER'S LIST

Slater [11] produced a list of Rogers-Ramanujan type series/product identities; Sills [10] in an amazing effort reworked and annotated this list, providing in particular finite versions of all of them.

Arguably the second most popular identities in the Rogers-Ramanujan world are Slater's [11] identities (39) and (38)

$$\begin{aligned} \sum_{n \geq 0} \frac{q^{2n^2}}{(q; q)_{2n}} &= \prod_{k \geq 1, k \equiv \pm 2, \pm 3, \pm 4, \pm 5 \pmod{16}} \frac{1}{1-q^k}, \\ \sum_{n \geq 0} \frac{q^{2n^2+2n}}{(q; q)_{2n+1}} &= \prod_{k \geq 1, k \equiv \pm 1, \pm 4, \pm 6, \pm 7 \pmod{16}} \frac{1}{1-q^k}. \end{aligned}$$

Let

$$s_n(z) = \sum_{0 \leq 2h \leq n} q^{2h^2} \begin{bmatrix} n \\ 2h \end{bmatrix} z^h \quad \text{and} \quad t_n(z) = \sum_{0 \leq 2h \leq n} q^{2h^2+2h} \begin{bmatrix} n \\ 2h+1 \end{bmatrix} z^h;$$

these polynomials are called *Santos* polynomials [4, 2, 3].

In order to describe the finite continued fraction expansion of $zt_n(z)/s_n(z)$, we define the following numbers and polynomials (power series) which were originally found by guessing:

$$a_{2k} := \frac{(1 - q^{4k+1})(q^{n+1-2k}; q^2)_{2k}}{q^{2k}(q^{n-2k}; q^2)_{2k+1}},$$

$$a_{2k+1} := \frac{(1 - q^{4k+3})(q^{n-2k}; q^2)_{2k+1}}{q^{2k+2}(q^{n-2k-1}; q^2)_{2k+2}};$$

$$S_{2i} := \sum_{j \geq 0} \frac{q^{2(i+j)(i+j+1)}(q^{n-2i-2j}; q)_{2j}(q^{n-2i}; q^2)_{2i+1}}{(q; q)_{2j+1}(q^{2j+3}; q^2)_{2i}} z^j,$$

$$S_{2i+1} := \sum_{j \geq 0} \frac{q^{2(i+j+1)^2}(q^{n-1-2i-2j}; q)_{2j}(q^{n-1-2i}; q^2)_{2i+2}}{(q; q)_{2j+1}(q^{2j+3}; q^2)_{2i+1}} z^j.$$

Then $zS_{i+1} = S_{i-1} - a_i S_i$, $S_{-1} = s_n(z)$, $S_0 = t_n(z)$.

We note that for $n \rightarrow \infty$, the quantities a_i and S_i appear already in [9].

The recursion will be shown for even i , the other instance being very similar:

$$\begin{aligned} S_{2i-1} - a_{2i} S_{2i} &= \sum_{j \geq 0} \frac{q^{2(i+j)^2}(q^{n+1-2i-2j}; q)_{2j}(q^{n+1-2i}; q^2)_{2i}}{(q; q)_{2j+1}(q^{2j+3}; q^2)_{2i-1}} z^j \\ &\quad - \frac{(1 - q^{4i+1})(q^{n+1-2i}; q^2)_{2i}}{q^{2i}(q^{n-2i}; q^2)_{2i+1}} \sum_{j \geq 0} \frac{q^{2(i+j)(i+j+1)}(q^{n-2i-2j}; q)_{2j}(q^{n-2i}; q^2)_{2i+1}}{(q; q)_{2j+1}(q^{2j+3}; q^2)_{2i}} z^j \\ &= (q^{n+1-2i}; q^2)_{2i} \sum_{j \geq 0} \frac{q^{2(i+j)^2}(q^{n+1-2i-2j}; q)_{2j}}{(q; q)_{2j+1}(q^{2j+3}; q^2)_{2i-1}} z^j \\ &\quad - (1 - q^{4i+1})(q^{n+1-2i}; q^2)_{2i} \sum_{j \geq 0} \frac{q^{2(i+j)^2+2j}(q^{n-2i-2j}; q)_{2j}}{(q; q)_{2j+1}(q^{2j+3}; q^2)_{2i}} z^j \\ &= (q^{n+1-2i}; q^2)_{2i} \sum_{j \geq 0} \frac{q^{2(i+j)^2}(q^{n+1-2i-2j}; q)_{2j-1}(1 - q^{n-2i})(1 - q^{2j+4i+1})}{(q; q)_{2j+1}(q^{2j+3}; q^2)_{2i}} z^j \\ &\quad - (1 - q^{4i+1})(q^{n+1-2i}; q^2)_{2i} \sum_{j \geq 0} \frac{q^{2(i+j)^2+2j}(q^{n+1-2i-2j}; q)_{2j-1}(1 - q^{n-2i-2j})}{(q; q)_{2j+1}(q^{2j+3}; q^2)_{2i}} z^j \\ &= (q^{n+1-2i}; q^2)_{2i} \sum_{j \geq 0} \frac{q^{2(i+j)^2}(q^{n+1-2i-2j}; q)_{2j-1}}{(q; q)_{2j+1}(q^{2j+3}; q^2)_{2i}} z^j \\ &\quad \times \left[(1 - q^{n-2i})(1 - q^{2j+4i+1}) - q^{2j}(1 - q^{4i+1})(1 - q^{n-2i-2j}) \right] \end{aligned}$$

$$\begin{aligned}
&= (q^{n+1-2i}; q^2)_{2i} \sum_{j \geq 0} \frac{q^{2(i+j)^2} (q^{n+1-2i-2j}; q)_{2j-1}}{(q; q)_{2j+1} (q^{2j+3}; q^2)_{2i}} z^j (1 - q^{2j}) (1 - q^{n+1+2i}) \\
&= (q^{n+1-2i}; q^2)_{2i+1} \sum_{j \geq 1} \frac{q^{2(i+j)^2} (q^{n+1-2i-2j}; q)_{2j-1}}{(q; q)_{2j+1} (q^{2j+3}; q^2)_{2i}} z^j (1 - q^{2j}) \\
&= z (q^{n-1-2i}; q^2)_{2i+2} \sum_{j \geq 0} \frac{q^{2(i+j+1)^2} (q^{n-1-2i-2j}; q)_{2j}}{(q; q)_{2j+1} (q^{2j+3}; q^2)_{2i+1}} z^j = z S_{2i+1}.
\end{aligned}$$

Further,

$$S_{-1} = \sum_{j \geq 0} \frac{q^{2j^2} (q^{n+1-2j}; q)_{2j}}{(q; q)_{2j+1} (q^{2j+3}; q^2)_{-1}} z^j = \sum_{j \geq 0} \frac{q^{2j^2} (q^{n+1-2j}; q)_{2j}}{(q; q)_{2j}} z^j = s_0(z)$$

and

$$S_0 = \sum_{j \geq 0} \frac{q^{2j(j+1)} (q^{n-2j}; q)_{2j} (1 - q^n)}{(q; q)_{2j+1}} z^j = \sum_{j \geq 0} \frac{q^{2j(j+1)} (q^{n-2j}; q)_{2j+1}}{(q; q)_{2j+1}} z^j = t_0(z).$$

Now we can iterate this relation in the following form:

$$\frac{zt(n)}{s(n)} = \frac{zS_0}{S_{-1}} = \frac{z}{a_0 + \frac{zS_1}{S_0}} = \frac{z}{a_0 + \frac{z}{a_1 + \frac{zS_2}{S_1}}} = \dots$$

This is the desired (finite) continued fraction expansion. It can be written more elegantly as

$$\frac{zt(n)}{s(n)} = \frac{zb_0}{1 + \frac{zb_1}{1 + \frac{zb_2}{1 + \dots}}}$$

with

$$b_0 = \frac{1}{a_0} = \frac{1 - q^n}{1 - q} \quad \text{and} \quad b_i = \frac{1}{a_{i-1} a_i} = \frac{q^{2i} (1 - q^{n-i}) (1 - q^{n+i})}{(1 - q^{2i-1}) (1 - q^{2i+1})} \quad \text{for } i \geq 1.$$

4. CONCLUSION

We would like to mention that it is more challenging to find the continued fractions and the relevant quantities, as the present proofs (and possibly other ones) consist of routine manipulations.

Since there are many Rogers-Ramanujan type identities and polynomials approximating them are not even unique, there should a plethora of additional results; even our previous effort [9] was 45 pages long!

Keeping this in mind, this paper is an invitation to young people with stamina to find many of these finite continued fractions!

REFERENCES

- [1] George E. Andrews. *The theory of partitions*. Addison-Wesley Publishing Co., Reading, Mass.-London-Amsterdam, 1976. Encyclopedia of Mathematics and its Applications, Vol. 2.
- [2] George E. Andrews, Arnold Knopfmacher, and Peter Paule. An infinite family of Engel expansions of Rogers-Ramanujan type. *Adv. in Appl. Math.*, 25(1):2–11, 2000.
- [3] George E. Andrews, Arnold Knopfmacher, Peter Paule, and Helmut Prodinger. q -Engel series expansions and Slater's identities. *Quaest. Math.*, 24(3):403–416, 2001. Dedicated to the memory of John Knopfmacher.
- [4] George E. Andrews and J. Plínio O. Santos. Rogers-Ramanujan type identities for partitions with attached odd parts. *Ramanujan J.*, 1(1):91–99, 1997.
- [5] Bruce C. Berndt. *Ramanujan's notebooks. Part III*. Springer-Verlag, New York, 1991.
- [6] Bruce C. Berndt, Sen-Shan Huang, Jaebum Sohn, and Seung Hwan Son. Some theorems on the Rogers-Ramanujan continued fraction in Ramanujan's lost notebook. *Trans. Amer. Math. Soc.*, 352(5):2157–2177, 2000.
- [7] D. M. Bressoud. Some identities for terminating q -series. *Math. Proc. Cambridge Philos. Soc.*, 89(2):211–223, 1981.
- [8] Robin Chapman. A new proof of some identities of Bressoud. *Int. J. Math. Math. Sci.*, 32:627–633, 2002.
- [9] Nancy S. S. Gu and Helmut Prodinger. On some continued fraction expansions of the Rogers-Ramanujan type. *Ramanujan J.*, 26(3):323–367, 2011.
- [10] Andrew V. Sills. Finite Rogers-Ramanujan type identities. *Electron. J. Combin.*, 10:Research Paper 13, 122, 2003.
- [11] L. J. Slater. Further identities of the Rogers-Ramanujan type. *Proc. London Math. Soc. (2)*, 54:147–167, 1952.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF STELLENBOSCH 7602, STELLENBOSCH, SOUTH AFRICA
E-mail address: hprodinger@sun.ac.za