Fountains, histograms, and q–identities

Peter Paule[1†] and Helmut Prodinger[2‡]

[1] *Research Institute for Symbolic Computation, Johannes Kepler University Linz, A-4040 Linz, Austria*
e-mail: Peter.Paule@risc.uni-linz.ac.at
Url: http://www.risc.uni-linz.ac.at/research/combinat/

[2] *The John Knopfmacher Centre for Applicable Analysis and Number Theory, School of Mathematics, University of the Witwatersrand, P. O. Wits, 2050 Johannesburg, South Africa*
e-mail: helmut@maths.wits.ac.za
Url: http://www.wits.ac.za/helmut/

received —, revised —, accepted —.

We solve the recursion $S_n = S_{n-1} - q^n S_{n-p}$, both, explicitly, and in the limit for $n \to \infty$, proving in this way a formula due to Merlini and Sprugnoli. It is also discussed how computer algebra could be applied.

Keywords: q–identities, fountains, histograms, Schur polynomials

1 Fountains and histograms

Merlini and Sprugnoli [6] discuss *fountains* and *histograms*; for the reader's convenience, we review a few key issues here.

A *fountain with n coins* is an arrangement of n coins in rows such that each coin in a higher row touches exactly two coins in the next lower row.

A *p-histogram* is a sequence of columns in which the height of the $(j+1)$st column is at most $k+p$, if k is the height of column j; the first column has height r, with $1 \le r \le p$.

It can be shown that the enumeration of coins in a fountain is equivalent with the enumeration of 1-histograms. The paper [6] addresses the enumeration of p-histograms with respect to area (=number of cells). Let $f_n^{[p]}$ be the number p-histograms with area n and $F^{[p]}(q)$ the corresponding generating function $F^{[p]}(q) = \sum_n f_n^{[p]} q^n$. The authors of [6] use two different approaches: one produces the answer in the form

$$F^{[p]}(q) = \lim_{m\to\infty} \frac{D_m}{E_m},$$

†Partially supported by SFB grant F1305 of the Austrian FWF.
‡The research was started when this author visited the RISC in 2002. Partially supported by NRF grant 2053748.

1365–8050 © Maison de l'Informatique et des Mathématiques Discrètes (MIMD), Paris, France

with some polynomials D_m, E_m defined in the next section, and the other gives it as

$$F^{[p]}(q) = \sum_{k\ge 0} \frac{(-1)^k q^{p\binom{k+1}{2}}}{(1-q)\dots(1-q^k)} \Big/ \sum_{k\ge 0} \frac{(-1)^k q^{k+p\binom{k}{2}}}{(1-q)\dots(1-q^k)}.$$

According to [5], it would be nice to have a direct argument that these two answers coincide. This is the subject of the present note.

2 Generalized Schur polynomials

The polynomials mentioned in the introduction are for fixed $p \ge 1$ defined as follows:

$$E_n = E_{n-1} - q^n E_{n-p}, \quad n \ge p, \qquad E_0 = \dots = E_{p-1} = 1,$$
$$D_n = D_{n-1} - q^n D_{n-p}, \quad n \ge p, \qquad D_i = 1 - \sum_{j=1}^{i} q^j,\ i = 0, \dots, p-1.$$

They can be compared with the classical Schur polynomials [8], which occur for $p = 2$ and $q = -1$. Then Merlini and Sprugnoli want a direct proof of the formulæ

$$E_\infty := \lim_{n\to\infty} E_n = \sum_{k\ge 0} \frac{(-1)^k q^{p\binom{k+1}{2}}}{(1-q)\dots(1-q^k)},$$
$$D_\infty := \lim_{n\to\infty} D_n = \sum_{k\ge 0} \frac{(-1)^k q^{k+p\binom{k}{2}}}{(1-q)\dots(1-q^k)}.$$

We will not only achieve that but actually derive *explicit* expressions for these polynomials!

It should be mentioned that Cigler [4] developed independently a combinatorial method to deal with recursions as ours, but also more general ones.

Let us study the generic recursion

$$S_n = S_{n-1} + t q^{n-p} S_{n-p},$$

with unspecified initial values $S_0, \dots, S_{p-1}$. For $p = 2$, these polynomials were studied by Andrews (and others) in the context of *Schur polynomials*, see [2].

We will use standard notation from q–calculus, see [1]:

$$(x)_n = (1-x)(1-xq)\dots(1-xq^{n-1}), \qquad \begin{bmatrix} n \\ k \end{bmatrix} = \frac{(q)_n}{(q)_k (q)_{n-k}}.$$

It will be convenient to define $\begin{bmatrix} n \\ k \end{bmatrix} = 0$ for $n < 0$ or $k > n$.

Now we will proceed as in [1] and consider noncommutative variables x, η, such that $x\eta = q\eta x$; all other variables commute.

Lemma 1.

$$\left(x + x^p \eta\right)^n = \sum_{k=0}^{n} \begin{bmatrix} n \\ k \end{bmatrix} q^{p\binom{n}{2} - pnk + p\binom{k+1}{2}} x^{k+p(n-k)} \eta^{n-k}.$$

Proof. We write

$$\left(x+x^p\eta\right)^n = \sum_{k=0}^{n} a_{n,k} x^{k+p(n-k)} \eta^{n-k},$$

and $\left(x+x^p\eta\right)^{n+1} = \left(x+x^p\eta\right)^n \left(x+x^p\eta\right)$ resp. as $\left(x+x^p\eta\right)^{n+1} = \left(x+x^p\eta\right)\left(x+x^p\eta\right)^n$, compare coefficients, and get the recursions

$$\begin{aligned} a_{n+1,k} &= a_{n,k-1} + a_{n,k} q^{k+p(n-k)}, \\ a_{n+1,k} &= a_{n,k-1} q^{n+1-k} + a_{n,k} q^{p(n-k)}. \end{aligned}$$

From this we derive, taking differences,

$$a_{n,k} = \frac{1-q^{n+1-k}}{1-q^k} q^{-p(n-k)} a_{n,k-1}.$$

The result follows from iteration by noting that $a_{n,0} = q^{p\binom{n}{2}}$. □

Of course we also have

$$\left(x+tx^p\eta\right)^n = \sum_{k=0}^{n} \begin{bmatrix} n \\ k \end{bmatrix} q^{p\binom{n}{2} - pnk + p\binom{k+1}{2}} x^{k+p(n-k)} t^{n-k} \eta^{n-k}.$$

Now we derive the generating function for

$$F(x) = \sum_{n \geq 0} S_n x^n;$$

the following procedure is inspired by [2]. Note that we can alternatively view η as an operator, defined by $\eta f(x) = f(qx)$. Cigler worked also much with this technique [3, 4]. We find

$$\sum_{n \geq p} S_n x^n = \sum_{n \geq p} S_{n-1} x^n + \sum_{n \geq p} t q^{n-p} S_{n-p} x^n = x \sum_{n \geq p-1} S_n x^n + t x^p \sum_{n \geq 0} \eta S_n x^n$$

or

$$F(x) - \sum_{n<p} S_n x^n = xF(x) - x \sum_{n<p-1} S_n x^n + t x^p \eta F(x),$$

and

$$F(x) = \frac{1}{1-x-tx^p\eta} \left(\sum_{i<p} S_i x^i - \sum_{i<p-1} S_i x^{i+1} \right).$$

Now we can apply our lemma and write

$$\begin{aligned} F(x) &= \sum_{n \geq 0} \left(x+tx^p\eta\right)^n \left(\sum_{i<p} S_i x^i - \sum_{i<p-1} S_i x^{i+1} \right) \\ &= \sum_{n \geq 0} \sum_{k=0}^{n} \begin{bmatrix} n \\ k \end{bmatrix} q^{p\binom{n}{2} - pnk + p\binom{k+1}{2}} x^{k+p(n-k)} t^{n-k} \eta^{n-k} \left(\sum_{i<p} S_i x^i - \sum_{i<p-1} S_i x^{i+1} \right) \\ &= \sum_{n \geq 0} \sum_{k=0}^{n} \begin{bmatrix} n \\ k \end{bmatrix} q^{p\binom{n}{2} - pnk + p\binom{k+1}{2}} x^{k+p(n-k)} t^{n-k} \left(\sum_{i<p} S_i q^{i(n-k)} x^i - \sum_{i<p-1} S_i q^{(i+1)(n-k)} x^{i+1} \right) \end{aligned}$$

$$= \sum_{n\geq 0}\sum_{k=0}^{n} \begin{bmatrix} n \\ k \end{bmatrix} q^{p\binom{k}{2}} x^{n-k+pk} t^k \left(\sum_{i<p} S_i q^{ik} x^i - \sum_{i<p-1} S_i q^{(i+1)k} x^{i+1} \right)$$

$$= \sum_{k,n\geq 0} \begin{bmatrix} n+k \\ k \end{bmatrix} q^{p\binom{k}{2}} x^{n+pk} t^k \left(\sum_{i<p} S_i q^{ik} x^i - \sum_{i<p-1} S_i q^{(i+1)k} x^{i+1} \right)$$

$$= \sum_{k\geq 0} q^{p\binom{k}{2}} x^{pk} t^k \frac{1}{(x)_{k+1}} \left(\sum_{i<p} S_i q^{ik} x^i - \sum_{i<p-1} S_i q^{(i+1)k} x^{i+1} \right).$$

From this we find an explicit formula for S_n (the quantity S_{-1} has to be interpreted as 0):

$$S_n = \sum_{0\leq i<p} (S_i - S_{i-1}) \sum_{k\geq 0} \begin{bmatrix} n-(p-1)k-i \\ k \end{bmatrix} q^{p\binom{k}{2}+ik} t^k.$$

Now we specialize this to our instance. Here, $t = -q^p$, and thus

$$S_n = \sum_{0\leq i<p} (S_i - S_{i-1}) \sum_{k\geq 0} \begin{bmatrix} n-(p-1)k-i \\ k \end{bmatrix} q^{p\binom{k+1}{2}+ik} (-1)^k.$$

Therefore

$$E_n = \sum_{k\geq 0} \begin{bmatrix} n-(p-1)k \\ k \end{bmatrix} q^{p\binom{k+1}{2}} (-1)^k.$$

From this, the limit of E_n is immediate. For D_n we eventually get the following form

$$D_n = \sum_{k\geq 0} \begin{bmatrix} n-(p-1)(k-1) \\ k \end{bmatrix} q^{k+p\binom{k}{2}} (-1)^k,$$

from which the formula for D_∞ is immediate. To prove it, we need a simple lemma whose proof is just a routine calculation.

Lemma 2.

$$\begin{bmatrix} m-i \\ k \end{bmatrix} q^{i(k+1)} = g(i) - g(i-1) \qquad \text{where} \qquad g(i) = -\begin{bmatrix} m-i \\ k+1 \end{bmatrix} q^{(i+1)(k+1)}. \quad \square$$

Now we can plug into the general formula above and compute

$$\begin{aligned} D_n &= E_n - \sum_{i=1}^{p-1} \sum_{k\geq 0} \begin{bmatrix} n-(p-1)k-i \\ k \end{bmatrix} q^{p\binom{k+1}{2}+i(k+1)} (-1)^k \\ &= E_n - \sum_{k\geq 0} (-1)^k q^{p\binom{k+1}{2}} \sum_{i=1}^{p-1} \begin{bmatrix} n-(p-1)k-i \\ k \end{bmatrix} q^{i(k+1)} \\ &= E_n - \sum_{k\geq 0} (-1)^k q^{p\binom{k+1}{2}} \left\{ q^{k+1} \begin{bmatrix} n-(p-1)k \\ k+1 \end{bmatrix} - q^{p(k+1)} \begin{bmatrix} n-(p-1)(k+1) \\ k+1 \end{bmatrix} \right\} \\ &= 1 - \sum_{k\geq 0} (-1)^k q^{p\binom{k+1}{2}} q^{k+1} \begin{bmatrix} n-(p-1)k \\ k+1 \end{bmatrix}, \end{aligned}$$

which is the announced formula after a simple change of variable. Note that in the penultimate step the telescoping property of the lemma has been used.

3 Computer algebra proofs

The polynomial families (E_n) and (D_n) give rise to the following study with respect to possible computer proofs. Let us take as input our sum representations of E_n and D_n:

$$\begin{aligned} E_n &= \sum_{k\geq 0} \begin{bmatrix} n-(p-1)k \\ k \end{bmatrix} q^{p\binom{k+1}{2}} (-1)^k, \\ D_n &= \sum_{k\geq 0} \begin{bmatrix} n-(p-1)(k-1) \\ k \end{bmatrix} q^{k+p\binom{k}{2}} (-1)^k. \end{aligned} \tag{3.1}$$

Then, if p is chosen as a specific positive integer, Riese's package qZeil [7] returns the recurrences $S_n = S_{n-1} - q^n S_{n-p}$ $(n \geq p)$ together with a certificate function Cert for independent verification. Despite the fact that for general "generic" integer parameter p there is no algorithm available, a general pattern can be easily guessed from running the algorithm for $p = 1$, $p = 2$, and $p = 3$, say.

For example, let $F(n,k)$ be the kth summand in our sum representation (3.1) of E_n, then the recurrence for E_n can be refined to the following statement.

Theorem 3.1. *For $n \geq p$ and $\delta_k f(n,k) = f(n,k) - f(n,k-1)$, we have*

$$F(n,k) - F(n-1,k) + q^n F(n-p,k) = \delta_k \mathsf{Cert}(n,k) F(n,k), \tag{3.2}$$

where

$$\mathsf{Cert}(n,k) = q^n \frac{(q^{n-p(k+1)+1})_p}{(q^{n-(p-1)(k+1)})_p}.$$

Proof. After dividing both sides of (3.2) by $F(n,k)$ the proof reduces to checking equality of rational functions. Namely, note that

$$\begin{aligned} \frac{F(n-1,k)}{F(n,k)} &= \frac{1-q^{n-pk}}{1-q^{n-(p-1)k}}, \\ \frac{F(n,k-1)}{F(n,k)} &= -\frac{q^{pk}}{1-q^k} \frac{(q^{n-pk+1})_p}{(q^{n-(p-1)k+1})_{p-1}}, \end{aligned}$$

and

$$\frac{F(n-p,k)}{F(n,k)} = q^{-n} \mathsf{Cert}(n,k). \quad \square$$

Analogously, there is a refined version of the recurrence for D_n. The certificate in this case is

$$\mathsf{Cert}(n,k) = q^n \frac{(q^{n-pk})_p}{(q^{n-(p-1)k})_p}.$$

Summarizing, with the sum representation for E_n and D_n in hand, the corresponding recurrences follow immediately by summing both sides of the computer recurrences (3.2) over all $k \geq 0$.

References

[1] G. E. Andrews, R. Askey, and R. Roy, *Special functions*, Encyclopedia of Mathematics and its Applications, vol. 71, Cambridge University Press, 1999.

[2] G. E. Andrews, *Fibonacci numbers and Rogers–Ramanujan identities*, The Fibonacci Quarterly, to appear (2003), 15 pp.

[3] J. Cigler, *Elementare q–Identitäten*, Séminaire Lotharingien de Combinatoire **B05a** (1981), 29 pp.

[4] J. Cigler, *Some algebraic aspects of Morse code sequences*, Discrete Mathematics and Theoretical Computer Science **6** (2003), 55–68.

[5] D. Merlini, *Private communication*, (2002).

[6] D. Merlini and R. Sprugnoli, *Fountains and histograms*, J. Algorithms **44** (2002), no. 1, 159–176.

[7] P. Paule and A. Riese, *A Mathematica q–analogue of Zeilberger's algorithm based on an algebraically motivated approach to q–hypergeometric telescoping*, Special functions, q–series and related topics (Toronto, ON, 1995), Fields Inst. Commun., vol. 14, Amer. Math. Soc., Providence, RI, 1997, pp. 179–210.

[8] I. Schur, *Ein Beitrag zur additiven Zahlentheorie und zur Theorie der Kettenbrüche*, S.-B. Preuss. Akad. Wiss. Phys.-Math. Kl., 1917, 302–321, reprinted in I. Schur, Gesammelte Abhandlungen, vol. 2, pp. 117–136, Springer, 1973.