


# ON THE ENUMERATION OF HOPPY'S WALKS

HELMUT PRODINGER

## 1. HOPPY WALKS

Deng and Mansour [1] introduce a rabbit named Hoppy and let him move according to certain rules. At that stage, we don't need to know the rules. Eventually, the enumeration problem is one about  $k$ -Dyck paths. The up-steps are  $(1, k)$  and the down-steps are  $(1, -1)$ .



The question is about the length of the sequence of down-steps printed in red. Or, phrased differently, how many  $k$ -Dyck paths end on level  $j$ , after  $m$  up-steps, the last step being an up-step. The recent paper [6] contains similar computations, although without the restriction that the last step must be an up-step.

Counting the number of up-steps is enough, since in total, there are  $m + km = (k + 1)m$  steps. The original description of Deng and Mansour is a reflection of this picture, with up-steps of size 1 and down-steps of size  $-k$ , but we prefer it as given here, since we are going to use the adding-a-new-slice method, see [2, 5]. A slice is here a run of down-steps, followed by an up-step. The first up-step is treated separately, and then  $m - 1$  new slices are added. We keep track of the level after each slice, using a variable  $u$ . The variable  $z$  is used to count the number of up-steps.

Deng and Mansour work out a formula which comprises  $O(m)$  terms. Our method leads only to a sum of  $O(j)$  terms.

The following substitution is essential for adding a new slice:

$$u^j \longrightarrow z \sum_{0 \leq h \leq j} u^{h+k} = \frac{zu^k}{1-u}(1-u^{j+1}).$$

Now let  $F_m(z, u)$  be the generating function according to  $m$  runs of down-steps. The substitution leads to

$$F_{m+1}(z, u) = \frac{zu^k}{1-u}F_m(z, 1) - \frac{zu^{k+1}}{1-u}F_m(z, u), \quad F_0(z, u) = zu^k.$$

Let  $F = \sum_{m \geq 0} F_m$ , then

$$F(z, u) = zu^k + \frac{zu^k}{1-u}F(z, 1) - \frac{zu^{k+1}}{1-u}F(z, u),$$

or

$$F(z, u) \frac{1-u+zu^{k+1}}{1-u} = zu^k + \frac{zu^k}{1-u}F(z, 1).$$

The equation  $1-u+zu^{k+1}=0$  is famous when enumerating  $(k+1)$ -ary trees. Its relevant combinatorial solution (also the only one being analytic at the origin) is

$$\bar{u} = \sum_{\ell \geq 0} \frac{1}{1+\ell(k+1)} \binom{1+\ell(k+1)}{\ell} z^\ell.$$

Since  $u-\bar{u}$  is a factor of the LHS, it must also be a factor of the RHS, and we can compute (by dividing out the factor  $(u-\bar{u})$ ) that

$$\frac{zu^k(1-u+F(z, 1))}{u-\bar{u}} = -zu^k.$$

Thus

$$F(z, u) = zu^k \frac{\bar{u}-u}{1-u+zu^{k+1}}.$$

The first factor has even a combinatorial interpretation, as a description of the first step of the path. It is also clear from this that the level reached is  $\geq k$  after each slice. We don't care about the factor  $zu^k$  anymore, as it produces only a simple shift. The main interest is now how to get to the coefficients of

$$\frac{\bar{u}-u}{1-u+zu^{k+1}}$$

in an efficient way. There is also the formula

$$1-u+zu^{k+1} = (\bar{u}-u) \left( 1 - z \frac{u^{k+1} - \bar{u}^{k+1}}{u - \bar{u}} \right),$$

but it does not seem to be useful here.

First we deal with the denominators

$$S_j := [u^j] \frac{1}{1 - u + zu^{k+1}} = \sum_{0 \leq m \leq j/k} (-1)^m \binom{j - km}{m} z^m.$$

One way to see this formula is to prove by induction that the sums  $S_j$  satisfy the recursion

$$S_j - S_{j-1} + zS_{j-k-1} = 0$$

and initial conditions  $S_0 = \dots = S_k = 1$ . In [6] such expressions also appear as determinants. Summarizing,

$$\frac{1}{1 - u + zu^{k+1}} = \sum_{m \geq 0} (-1)^m z^m \sum_{j \geq km} \binom{j - km}{m} u^j.$$

Now we read off coefficients:

$$\begin{aligned} & [u^j] \frac{\bar{u}}{1 - u + zu^{k+1}} \\ &= \sum_{0 \leq m \leq j/k} (-1)^m \binom{j - km}{m} z^m \sum_{\ell \geq 0} \frac{1}{1 + \ell(k+1)} \binom{1 + \ell(k+1)}{\ell} z^\ell \end{aligned}$$

and further

$$\begin{aligned} & [z^n][u^j] \frac{\bar{u}}{1 - u + zu^{k+1}} \\ &= \sum_{0 \leq m \leq j/k} (-1)^m \binom{j - km}{m} \frac{1}{1 + (n - m)(k+1)} \binom{1 + (n - m)(k+1)}{n - m}. \end{aligned}$$

The final answer to the Deng-Mansour enumeration (without the shift) is

$$\begin{aligned} & \sum_{0 \leq m \leq j/k} (-1)^m \binom{j - km}{m} \frac{1}{1 + (n - m)(k+1)} \binom{1 + (n - m)(k+1)}{n - m} \\ & \quad - (-1)^n \binom{j - 1 - kn}{n}. \end{aligned}$$

If one wants to take care of the factor  $zu^k$  as well, one needs to do the replacements  $n \rightarrow n+1$  and  $j \rightarrow j+k$  in the formula just derived. That enumerates then the  $k$ -Dyck paths ending at level  $j$  after  $n$  up-steps, where the last step is an up-step.

## 2. AN APPLICATION

The encyclopedia of integer sequences [4] has the sequences A334680, A334682, A334719, (with a reference to [3]) which is the total number of down-steps of the last down-run, for  $k = 2, 3, 4$ . So, if the path ends on level  $j$ , the contribution to the total is  $j$ .

All we have to do here is to differentiate

$$F(z, u) = zu^k \frac{\bar{u} - u}{1 - u + zu^{k+1}}.$$

w.r.t.  $u$ , and then replace  $u$  by 1. The result is

$$\frac{\bar{u}}{z} - \bar{u} - \frac{1}{z},$$

and the coefficient of  $z^m$  therein is


$$\frac{1}{1 + (m+1)(k+1)} \binom{1 + (m+1)(k+1)}{m+1} - \frac{1}{1 + m(k+1)} \binom{1 + m(k+1)}{m}.$$

I don't know how this was derived in [3], but it is more fun to figure out things for oneself!

We hope to report about more applications soon.

## 3. HOPPY'S EARLY ADVENTURES

Now we investigate what Hoppy does after his first up-step; he might follow with  $0, 1, \dots, k$  down-steps. Eventually, we want to sum all these steps (red in the picture).



A new slice is now an up-step, followed by a sequence of down-steps. The substitution of interest is:

$$u^i \rightarrow z \sum_{0 \leq h \leq i+k} u^h = \frac{z}{1-u} - \frac{zu^{i+k+1}}{1-u}.$$

Furthermore

$$F_{h+1}(z, u) = \frac{z}{1-u} F_h(z, 1) - \frac{zu^{k+1}}{1-u} F_h(z, u),$$

and  $F_0 = u^h$ , the starting level.

We have

$$H(z, u) = \sum_{h \geq 0} F_h(z, u) = u^h + \frac{z}{1-u} H(z, 1) - \frac{zu^{k+1}}{1-u} H(z, u)$$

or

$$H(z, u)(1 - u + zu^{k+1}) = u^h(1 - u) + zH(z, 1)$$

Plugging in  $\bar{u}$  into the RHS gives 0:

$$zH(z, 1) = -\bar{u}^h(1 - \bar{u}),$$

and

$$H(z, u) = \frac{u^h(1 - u) - \bar{u}^h(1 - \bar{u})}{1 - u + zu^{k+1}}.$$

But we only need  $H(z, 0)$ , since we return to the  $x$ -axis at the end:

$$H(z, 0) = [h = 0] + \bar{u}^{h+1} - \bar{u}^h.$$

The total contribution of red steps is then

$$k + \sum_{h=0}^k (k - h)(\bar{u}^{h+1} - \bar{u}^h) = \sum_{h=1}^k \bar{u}^h;$$

the coefficient of  $z^m$  in this is the total contribution. Since  $\bar{u} = 1 + z\bar{u}^{k+1}$ , there is the further simplification

$$-1 + \frac{1}{z} + \frac{1}{1 - \bar{u}} = \sum_{m \geq 1} \frac{k}{m+1} \binom{(k+1)m}{m} z^m.$$

The proof of this is as follows. Let  $m \geq 1$ , then

$$\begin{aligned} [z^m] \left( -1 + \frac{1}{z} + \frac{1}{1 - \bar{u}} \right) &= -[z^m] \frac{1}{z\bar{u}^{k+1}} \\ &= -[z^{m+1}] \sum_{\ell \geq 0} \frac{-(k+1)}{(k+1)\ell - (k+1)} \binom{(k+1)\ell - (k+1)}{\ell} z^\ell \\ &= [z^{m+1}] \sum_{\ell \geq 0} \frac{(k+1)}{(k+1)(\ell - 1)} \binom{(k+1)(\ell - 1)}{\ell} z^\ell \\ &= \frac{(k+1)}{(k+1)m} \binom{(k+1)m}{m+1} = \frac{k}{m+1} \binom{(k+1)m}{m}. \end{aligned}$$

We did not expect such a simple answer  $\frac{k}{m+1} \binom{(k+1)m}{m}$  to this question about Hoppy's early adventures!

This analysis of Hoppy's early adventures covers sequences A007226, A007228, A124724 of [4], with references to [3].

#### REFERENCES

- [1] Eva Deng and Toufik Mansour, Three Hoppy path problems and ternary paths, *Discrete Applied Mathematics* 156, 2008, 770–779.
- [2] Flajolet, Philippe and Prodinger, Helmut, Level number sequences for trees, *Discrete Mathematics* 65, 1987, 149–156.
- [3] Andrei Asinowski, Benjamin Hackl, Sarah J. Selkirk, Down-step statistics in generalized Dyck path, arXiv:2007.15562, 2020.
- [4] The online encyclopedia of integer sequences. <http://oeis.org>.
- [5] Helmut Prodinger. Analytic methods. In *Handbook of enumerative combinatorics*, *Discrete Math. Appl.* (Boca Raton), pages 173–252. CRC Press, Boca Raton, FL, 2015.
- [6] Prodinger, Helmut, On  $k$ -Dyck paths with a negative boundary.

HELMUT PRODINGER, DEPARTMENT OF MATHEMATICAL SCIENCES, STELLENBOSCH UNIVERSITY, 7602 STELLENBOSCH, SOUTH AFRICA

*Email address:* hprodinger@sun.ac.za