

# AN IDENTITY CONJECTURED BY LACASSE VIA THE TREE FUNCTION

HELMUT PRODINGER

ABSTRACT. A. Lacasse conjectured a combinatorial identity in his study of learning theory. Various people found independent proofs. Here is another one that is based on the study of the tree function, with links to Lambert's  $W$ -function and Ramanujan's  $Q$ -function. It is particularly short.

Let

$$\xi(n) = \sum_{k=0}^n \binom{n}{k} \left(\frac{k}{n}\right)^k \left(\frac{n-k}{n}\right)^{n-k},$$

$$\xi_2(n) = \sum_{k_1+k_2+k_3=n} \frac{n!}{k_1!k_2!k_3!} \left(\frac{k_1}{n}\right)^{k_1} \left(\frac{k_2}{n}\right)^{k_2} \left(\frac{k_3}{n}\right)^{k_3}.$$

Then Lacasse [3] conjectured that  $\xi_2(n) = \xi(n) + n$ . There are now three independent proofs [6, 1, 5]. Here, we want to shed additional light on the matter, by using the *tree function* (equivalent to Lambert's  $W$ -function [7]) and linking the enumeration to the celebrated  $Q$ -function of Ramanujan [2].

As Younsi has already pointed out, it is easier to work with

$$\alpha(n) = n^n \xi(n) = \sum_{k=0}^n \binom{n}{k} k^k (n-k)^{n-k},$$

$$\beta(n) = n^n \xi_2(n) = \sum_{k_1+k_2+k_3=n} \frac{n!}{k_1!k_2!k_3!} k_1^{k_1} k_2^{k_2} k_3^{k_3}.$$

The tree function [4]  $y(z)$  is defined by  $y = ze^y$  and possesses the expansion

$$y(z) = \sum_{n \geq 1} n^{n-1} \frac{z^n}{n!}.$$

Therefore

$$\sum_{n \geq 0} n^n \frac{z^n}{n!} = 1 + zy'(z) = 1 + \frac{y}{1-y} = \frac{1}{1-y}.$$

Consequently,

$$\alpha(n) = n! [z^n] \left(\frac{1}{1-y}\right)^2 \quad \text{and} \quad \beta(n) = n! [z^n] \left(\frac{1}{1-y}\right)^3.$$

---

The author was supported by an incentive grant of the NRF of South Africa.

Now we compute the coefficients of a general power via Cauchy's integral formula:

$$\begin{aligned}
[z^n] \left( \frac{1}{1-y} \right)^d &= \frac{1}{2\pi i} \oint \frac{dz}{z^{n+1}} \left( \frac{1}{1-y} \right)^d \\
&= \frac{1}{2\pi i} \oint \frac{dy(1-y)e^{-y}e^{y(n+1)}}{y^{n+1}} \left( \frac{1}{1-y} \right)^d \\
&= [y^n] \frac{e^{ny}}{(1-y)^{d-1}} \\
&= \sum_{k=0}^n \frac{n^{n-k}}{(n-k)!} \binom{k+d-2}{d-2}.
\end{aligned}$$

Therefore

$$\alpha(n) = \sum_{k=0}^n \frac{n!}{k!} n^k, \quad \beta(n) = \sum_{k=0}^n \frac{n!}{k!} (n+1-k)n^k,$$

and

$$\begin{aligned}
\beta(n) - \alpha(n) &= \sum_{k=0}^n \frac{n!}{k!} (n-k)n^k \\
&= \sum_{k=0}^n \frac{n!}{k!} n^{k+1} - \sum_{k=1}^n \frac{n!}{(k-1)!} n^k = n^{n+1},
\end{aligned}$$

as claimed.

Note that  $\alpha(n)$  is linked to Ramanujan's celebrated  $Q$ -function [2] via

$$\alpha(n) = n^n (1 + Q(n)).$$

#### REFERENCES

- [1] W. Y. C. Chen, J. F. F. Peng and H. R. L. Yang, Decomposition of triply rooted trees, arXiv:1212.6468v1.
- [2] P. Flajolet, P. Grabner, P. Kirschenhofer and H. Prodinger, On Ramanujan's  $Q(n)$ -function, Journal of Computational and Applied Mathematics, 58:103–116, 1995.
- [3] A. Lacasse, Bornes PAC-Bayes et algorithmes d'apprentissage, Ph. D. thesis, Université Laval, Quebec, 2010.
- [4] I. Goulden and D. Jackson, Combinatorial Enumeration, Wiley, 1983.
- [5] Y. Sun, Another simple proof of an identity conjectured by Lacasse, arXiv:1209.0824.
- [6] M. Younsi, Proof of a combinatorial conjecture coming from the PAC-Bayesian machine learning theory, arXiv:1301.0679.
- [7] R. Corless, G. Gonnet, D. Hare, D., D. Jeffrey, D., D. Knuth, On the Lambert  $W$ -function, Advances in Computational Mathematics 5: 329-359, 1996.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF STELLENBOSCH 7602, STELLENBOSCH, SOUTH AFRICA

*E-mail address:* hprodinger@sun.ac.za